

UNIQUENESS IN INVERSE DIFFRACTION GRATING PROBLEMS WITH INFINITELY MANY PLANE WAVES AT A FIXED FREQUENCY*

XIAOXU XU[†], GUANGHUI HU[‡], BO ZHANG[§], AND HAIWEN ZHANG[¶]

Abstract. This paper is concerned with uniqueness of solution of the inverse diffraction by problems by a periodic curve with Dirichlet boundary condition in two dimensions. It is proved that the periodic curve can be uniquely determined by the near-field measurement data corresponding to infinitely many incident plane waves with distinct directions at a fixed frequency. Our proof is based on Schiffer's idea which consists of two ingredients: (i) the total fields for incident plane waves with distinct directions are linearly independent, and (ii) for a fixed wave number there exist only finitely many linearly independent Dirichlet eigenfunctions in a bounded domain or in a closed waveguide under additional assumptions on the waveguide boundary. Based on the Rayleigh expansion, we prove that the phased near-field data can be uniquely determined by the phaseless near-field data in a bounded domain, with the exception of a finite set of incident angles. Such a phase retrieval result leads to a new uniqueness result for the inverse grating diffraction problem with phaseless near-field data at a fixed frequency. Since the incident direction determines the quasi-periodicity of the boundary value problem, our inverse issues are different from the existing results of [F. Hettlich and A. Kirsch, *Inverse Problems*, 13 (1997), pp. 351–361], where fixed-direction plane waves at multiple frequencies were considered.

Key words. uniqueness, inverse grating diffraction problem, Dirichlet boundary condition, phaseless data

MSC codes. 35R30, 78A46, 35B27

DOI. 10.1137/22M1480963

1. Introduction. Suppose a perfectly conducting grating is illuminated by an incident monochromatic plane wave in an isotropic homogeneous background medium. For simplicity it is assumed that the grating is periodic in one surface direction x_1 and independent of another surface direction x_3 . In the present paper, we restrict discussions to the TE (transverse electric) polarization case, where the three-dimensional

* Received by the editors February 28, 2022; accepted for publication (in revised form) October 10, 2022; published electronically February 27, 2023.

<https://doi.org/10.1137/22M1480963>

Funding: The work of the first author is supported by China Postdoctoral Science Foundation grant 2019TQ0023, the Young Talent Support Plan of Xi'an Jiaotong University, the Fundamental Research Funds for the Central Universities grant number xzy012022009, and National Natural Science Foundation of China grants 12122114 and 12201489. The work of the second author is supported by National Natural Science Foundation of China grant 12071236, Fundamental Research Funds for Central Universities in China grant 63213025, and Beijing Natural Science Foundation grant Z210001. The work of the fourth author is supported by National Natural Science Foundation of China grant 11871466 and the Youth Innovation Promotion Association CAS.

[†] School of Mathematics and Statistics, Xi'an Jiaotong University, Xi'an, Shaanxi, 710049, China (xuxiaoxu@xjtu.edu.cn).

[‡] Corresponding author School of Mathematical Sciences and LPMC, Nankai University, Tianjin 300071, China (ghhu@nankai.edu.cn).

[§] LSEC and Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China and School of Mathematical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China (b.zhang@amt.ac.cn).

[¶] Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China (zhanghaiwen@amss.ac.cn).




FIG. 1. Scattering by a periodic curve with Dirichlet boundary condition.

scattering problem governed by the Maxwell equations can be reduced to a two-dimensional diffraction problem modeled by the scalar Helmholtz equation over the x_1x_2 -plane. Accordingly, the perfect conductor boundary condition on the grating surface can be reduced to the Dirichlet boundary condition. This work is concerned with the inverse diffraction problem of recovering the periodic curve (i.e., the cross-section of the grating surface) with a Dirichlet boundary condition from phased and phaseless near-field data measured above the grating.

Denote by $\Gamma \subset \mathbb{R}^2$ a curve that is periodic in the x_1 -direction and bounded in the x_2 -direction and represents the cross-section of the grating surface in the x_1x_2 -plane. Let the incident field be a time-harmonic plane wave of the form $u^i(x)e^{-i\omega t}$, incited at the angular frequency $\omega > 0$, where the spatially dependent function u^i takes the form

$$(1.1) \quad u^i(x) = e^{ikx \cdot d} = e^{ikx_1 \sin \theta - ikx_2 \cos \theta}, \quad x = (x_1, x_2) \in \mathbb{R}^2.$$

Here the incident direction $d := (\sin \theta, -\cos \theta)$ is given in terms of the incident angle $\theta \in (-\pi/2, \pi/2)$, and $k := \omega/c$ is the wave number with $c > 0$ denoting the wave speed in the homogeneous background medium. In this paper we assume further that Γ satisfies one of the following regularity conditions:

- Condition (i). Γ is the graph of a 3-times continuously differentiable function;
- Condition (ii). Γ is an analytical curve.

Denote by $L > 0$ the period of Γ and by Ω the unbounded connected domain above Γ (cf. Figure 1). The wave propagation is then modeled by the Dirichlet boundary value problem for the Helmholtz equation,

$$(1.2) \quad \Delta u + k^2 u = 0 \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \Gamma,$$

where the total field $u = u^i + u^s$ is the sum of the incident field u^i and the scattered field u^s .

Set $\alpha = \alpha(k, \theta) := k \sin \theta$. Obviously, the incident field (1.1) is α -quasi-periodic in the sense that $e^{-i\alpha x_1} u^i(x)$ is L -periodic with respect to x_1 for all $x \in \Omega$. In view of the periodicity of the structure together with the form of the incident field, we require the total field u to be α -quasi-periodic, that is, $e^{-i\alpha x_1} u(x)$ is L -periodic with respect to x_1 for all $x \in \Omega$. This implies that

$$(1.3) \quad u(x_1 + nL, x_2) = u(x_1, x_2)e^{i\alpha nL} \quad \text{for any } n \in \mathbb{Z}.$$

The number $\alpha \in \mathbb{R}$ will be referred to as the phase shift of the solution. Since the domain Ω is unbounded in the x_2 -direction, a radiation condition needs to be imposed at infinity as $x_2 \rightarrow \infty$ to ensure the well-posedness of the diffraction problem. Precisely, we require the scattered field u^s to satisfy the Rayleigh expansion, that is, there exist Rayleigh coefficients $A_n \in \mathbb{C}$ ($n \in \mathbb{Z}$) depending on k, θ and Γ such that

$$(1.4) \quad u^s(x) = \sum_{n \in \mathbb{Z}} A_n e^{i\alpha_n x_1 + i\beta_n x_2}, \quad x \in U_h := \{x = (x_1, x_2) \in \mathbb{R}^2 : x_2 > h\},$$

where the parameters $\alpha_n \in \mathbb{R}$ and $\beta_n \in \mathbb{C}$ for $n \in \mathbb{Z}$ are defined by

$$(1.5) \quad \begin{aligned} \alpha_n &= \alpha_n(k, \theta, L) := \alpha + 2n\pi/L, \\ \beta_n &= \beta_n(k, \theta, L) := \begin{cases} \sqrt{k^2 - (\alpha_n)^2} & \text{if } |\alpha_n| \leq k, \\ i\sqrt{(\alpha_n)^2 - k^2} & \text{if } |\alpha_n| > k \end{cases} \end{aligned}$$

for any fixed $h > \max\{x_2 : x \in \Gamma\}$. We note that the series (1.4) is uniformly convergent and bounded in U_h (see Lemma 2.1). It consists of a finite number of propagating wave modes for $|\alpha_n| \leq k$ and infinitely many surface (evanescent) wave modes corresponding to $|\alpha_n| > k$. For notational convenience we rewrite the incident plane wave (1.1) as

$$(1.6) \quad u^i(x) = A_i e^{i\alpha_i x_1 + i\beta_i x_2},$$

where $A_i = A_i(k, \theta) := 1$, $\alpha_i = \alpha_i(k, \theta) := k \sin \theta$, $\beta_i = \beta_i(k, \theta) := -k \cos \theta$. Here, the symbol i denotes the index for the incident plane wave. We note that $\alpha_i = \alpha = \alpha_0$ and $\beta_i = -\beta_0$.

The well-posedness of the forward diffraction problem is presented in the following proposition.

PROPOSITION 1.1. (1) *If Condition (i) holds, the diffraction problem (1.1)–(1.4) admits a unique α -quasi-periodic solution $u \in C^2(\Omega) \cap C(\overline{\Omega})$.*

(2) *Under Condition (ii), there exists at least one solution $u \in C^2(\Omega) \cap C(\overline{\Omega})$ to the diffraction problem (1.1)–(1.4). Moreover, uniqueness of the solution remains true for small wave numbers or for all wave numbers excluding a discrete set with the only accumulating point at infinity.*

We refer the reader to [20, 28] for the proof of the first statement when the period of the curve is $L = 2\pi$. Actually, it follows from a scaling argument that statement (1) holds for an arbitrary period $L > 0$. Further, by the Fredholm alternative (see, e.g., [35, Theorem 4.17]) and the analytic Fredholm theory (see, e.g., [14, Theorem 8.26]), one can prove the second statement through a standard variational argument together with quasi-periodic transparent boundary conditions (see, e.g., [1, 4, 10, 41]). We remark that the well-posedness of the diffraction problem (1.1)–(1.4) can be established under weaker conditions than Conditions (i) and (ii). To be more specific, if Γ is the graph of a periodic Lipschitz function, the existence of α -quasi-periodic variational solutions in $H_{0,\alpha}^1(\Omega)$ can be shown, where

$$H_{0,\alpha}^1(\Omega) := \{u \in H_{loc}^1(\Omega) : e^{-i\alpha x_1} u(x) \text{ is } L\text{-periodic with respect to } x_1, u = 0 \text{ on } \Gamma\}.$$

Further, uniqueness of solutions remains valid for any $k > 0$ even under the following weaker assumption (see [12, (4.1) and Theorem 4.1] and [11, (2.2) and Theorem 4.1]):

$$(x_1, x_2) \in \Omega \Rightarrow (x_1, x_2 + s) \in \Omega \quad \text{for all } s > 0.$$

Note that this geometric assumption is fulfilled if Γ is the graph of a continuous function.

The inverse problem we consider in this paper is to recover a periodic curve with Dirichlet boundary condition from phased or phaseless near-field data corresponding to an infinite number of incident plane waves with different angles, where the period L of the curve is unknown.

Let $\theta_n \in (-\pi/2, \pi/2)$ with $n \in \mathbb{Z}_+$ be distinct incident angles, and denote by $u(x; \theta_n)$ the total field corresponding to the diffraction problem (1.1)–(1.4) with $\theta = \theta_n$.

Note that, according to Proposition 1.1, the diffraction problem (1.1)–(1.4) may admit multiple solutions under Condition (ii) if k is an exceptional wave number. If this happens, $u(x; \theta_n)$ is assumed to be any one of these solutions. The main uniqueness result for the inverse problem considered is presented in the following theorem.

THEOREM 1.1. *Assume that the unknown periodic curve Γ with Dirichlet boundary condition satisfies either Condition (i) or Condition (ii). Suppose the period of Γ is unknown. Then Γ can be uniquely determined by either the phased data $\{u(x; \theta_n) : x \in \mathcal{S}\}_{n=1}^\infty$, where $\mathcal{S} \subset \Gamma_h$ is a line segment parallel to the x_1 -axis, or the phaseless data $\{|u(x; \theta_n)| : x \in \mathcal{D}\}_{n=1}^\infty$, where $\mathcal{D} \subset \Omega$ is a bounded domain. Here, $\Gamma_h := \{x : x_2 = h\}$ with $h > \max\{x_2 : x \in \Gamma\}$ being an arbitrary constant.*

The proof of Theorem 1.1 will be given in section 4 for the case of phased data and in section 5 for the case of phaseless data. If the background medium is nonabsorbing (i.e., $k > 0$), it is well known that the global uniqueness with phased near-field data corresponding to one incident plane wave is impossible (see [16]). We will show in section 3 that phased near-field data corresponding to one incident plane wave cannot even determine the period of a grating curve.

Below we shall review some uniqueness results for inverse time-harmonic scattering from periodic curves of Dirichlet kind and compare our results with them. To the best of our knowledge, uniqueness with one incident wave was verified in the following special cases:

- (i) when the background medium is lossy (i.e., $\text{Im } k > 0$) [6];
- (ii) when the wave number or the grating height is sufficiently small [23];
- (iii) within the class of rectangular gratings [19], or within the class of polygonal gratings when Rayleigh frequencies are excluded (i.e., $\beta_n \neq 0$ for all $n \in \mathbb{Z}$) [16].

If a Rayleigh frequency occurs (i.e., $\beta_n = 0$ for some $n \in \mathbb{Z}$), the measured data for two incident plane waves can be used to determine a general polygonal grating [19] (see also [8, 9] in the case of inverse electromagnetic scattering from perfectly conducting polyhedral gratings). Based on the “blowing up” arguments arising from inverse scattering from bounded obstacles [25, 30], Kirsch proved in [29] that a general periodic curve can be uniquely determined by using all α -quasi-periodic incident waves $\{e^{i\alpha_n x_1 - i\beta_n x_2} : n \in \mathbb{Z}\}$. The uniqueness result of [29] also extends to Neumann and impedance boundary value problems as well as the transmission problem [18, 24, 42]. Note that the incident waves used in [29] include a finite number of plane waves for $|\alpha_n| \leq k$ and infinitely many evanescent waves corresponding to $|\alpha_n| > k$, which constitute the α -quasi-periodic Green’s function with a fixed phase shift. However, the incoming evanescent waves are usually difficult to generate. The factorization method established in [5] also gives rise to the same uniqueness result. Our work differs from [5, 29] in the use of infinitely many plane waves with various directions lying on the semicircle $\mathbb{S}^- := \{x = (x_1, x_2) \in \mathbb{R}^2 : |x| = 1, x_2 < 0\}$. Such incoming waves do not include any evanescent wave and thus seem more natural in applications.

In the case when the a priori information of the grating height is available, Hettlich and Kirsch [23] obtained a uniqueness result by using fixed-direction plane waves with a finite number of frequencies. See also [21] for an analogous result for the inverse transmission problem in a two-layered periodic medium. These works can be viewed as the extension of the idea due to Colton and Sleeman [15] from inverse scattering by bounded sound-soft obstacles to the case of inverse scattering by periodic structures. As will be seen in subsection 4.2, the fixed-direction problem of [23] and the fixed-frequency problem to be investigated here are different inverse issues. With multiple incident directions, Colton and Sleeman’s idea [15] and the approach of

[23] cannot be straightforwardly applied to inverse diffraction problems in periodic structures, because of the different phase shifts caused by incident angles at a fixed frequency. In fact, the multifrequency inverse problem of [23] and the multidirection problem of this paper result in different eigenvalue problems. Using different directions leads to a μ -eigenvalue problem where $\mu = \sin \theta$ is determined by the incident angle $\theta \in (-\pi/2, \pi/2)$, which brings difficulties in proving the discreteness of eigenvalues. To apply the analytical Fredholm theory, we shall resort to the arguments of [40] to exclude the existence of flat dispersion curves in a closed waveguide. For this reason we need the same regularity assumption on grating profiles as in [40] (see Conditions (i) and (ii)).

In many practical applications, it is difficult to accurately measure the phase information of wave fields. This motivates us to study the inverse problem of recovering a periodic curve of Dirichlet kind from phaseless data. However, most uniqueness results with phaseless data are confined to inverse scattering from bounded scatterers (see, e.g., [26, 31, 32, 33, 38, 45]). In particular, using the Sommerfeld radiating behavior of the scattered field at infinity, we see that explicit formulas for recovering phased far-field patterns from phaseless near-field data are derived in [38]. This result has been extended to scattering by locally rough surfaces in [44]. The key ingredient of [38, 44] is the Rellich's lemma (see [14, Theorem 2.14] and [43, Theorem 2.2]), which establishes the one-to-one correspondence between Sommerfeld radiating waves and their far-field patterns. In this paper we consider a periodic curve (which can be viewed as a special nonlocally perturbed rough surface in two dimensions) and also prove a phase retrieval result but rather based on the Rayleigh expansion (1.4) for diffraction grating problems. We remark that the Rayleigh expansion radiation condition is different from the Sommerfeld radiation condition, since the scattered field (1.4) does not decay as x_2 tends to infinity and does not lead to the classical far-field patterns. As a consequence, the phase retrieval approach and uniqueness proofs in [38, 44] cannot carry over to diffraction grating problems here. Instead of recovering the phased far-field pattern as in [38, 44], we need to determine the Rayleigh coefficients in a special order by sophisticated arguments (see the proof of Theorem 5.2). To the best of our knowledge, uniqueness results for identifying periodic grating curves using phaseless near-field data are not yet available. We refer the reader to [2, 3, 7, 26, 34, 44, 46, 47] for numerical schemes of inverse scattering using phaseless data.

This paper is organized as follows. In section 2, we prepare several lemmas for later use. Section 3 is devoted to determining one grating period from the phased near-field data for one incident plane wave. The results in sections 2 and 3 are independent of the smoothness Conditions (i) and (ii) of the periodic curve given in the introduction. In section 4, we prove uniqueness for recovering periodic curves with Dirichlet boundary condition using the phased near-field data corresponding to infinitely many incident plane waves with distinct directions. A similar uniqueness result based on phaseless near-field data will be established in section 5. Finally, concluding remarks will be given in section 6.

2. Preliminary lemmas. The following lemmas are useful in the proofs of uniqueness results in what follows.

LEMMA 2.1. *Let Γ be a periodic curve. Set $U_h := \{x \in \mathbb{R}^2 : x_2 > h\}$ for any $h > \max\{x_2 : x \in \Gamma\}$.*

- (i) *The Rayleigh expansion (1.4) is uniformly bounded for $x \in U_h$.*
- (ii) *The Rayleigh expansion (1.4) is uniformly and absolutely convergent for $x \in U_h$.*

(iii) Let $b \in \mathbb{R}$, and let A_n ($n \in \mathbb{Z}$) be given as in (1.4). Set $\mathcal{P}_\pm(N) := \{n \in \mathbb{Z} : |\alpha_n| > k, \pm n > N\}$ for $N > 0$. Then, for the case when $b < |\beta_n|$ for all $n \in \mathcal{P}_+(N)$, the series

$$(2.1) \quad \sum_{n \in \mathcal{P}_+(N)} A_n e^{i\alpha_n x_1 + i\beta_n x_2 + b x_2}$$

is uniformly and absolutely convergent for $x \in U_h$. For the case when $b < |\beta_n|$ for all $n \in \mathcal{P}_-(N)$, the series

$$\sum_{n \in \mathcal{P}_-(N)} A_n e^{i\alpha_n x_1 + i\beta_n x_2 + b x_2}$$

is uniformly and absolutely convergent for $x \in U_h$.

(iv) Let $N \in \mathbb{Z}_+$, $a_j \in \mathbb{C}$, and $b_j \in \mathbb{R} \setminus \{0\}$ for $j = 1, \dots, N$. Then

$$\left| \frac{1}{T} \int_T^{2T} \sum_{j=1}^N a_j e^{ib_j t} dt \right| \leq \frac{2}{T} \sum_{j=1}^N \frac{|a_j|}{|b_j|} \rightarrow 0 \quad \text{as } T \rightarrow +\infty.$$

Proof. Choosing $\sigma > 0$ small enough so that $h - 2\sigma > \max\{x_2 : x \in \Gamma\}$, noting that (1.4) also holds with h replaced by $h - 2\sigma$, and applying Parseval's equality yield the estimate

$$(2.2) \quad \begin{aligned} 2|u^s(x)| &\leq \sum_{n \in \mathbb{Z}} 2|A_n e^{i\alpha_n x_1 + i\beta_n x_2}| \\ &\leq \sum_{n \in \mathbb{Z}} \left| A_n e^{i\alpha_n x_1 + i\beta_n (h-\sigma)} \right|^2 + \sum_{n \in \mathbb{Z}} \left| e^{i\beta_n (x_2 - h + \sigma)} \right|^2 \\ &\leq \frac{1}{L} \int_0^L |u^s(x_1, h - \sigma)|^2 dx_1 + \sum_{|\alpha_n| \leq k} 1 + \sum_{|\alpha_n| > k} C e^{-|n|/C} \end{aligned}$$

uniformly for all $x \in U_h$, where we have used the fact that $2\sigma \sqrt{(2n\pi/L + \alpha_0)^2 - k^2} > |n|/C$ holds for sufficiently large $|n|$ provided the constant $C > 0$ is large enough. Thus statement (i) holds. The estimate (2.2) also implies that statement (ii) holds.

We now prove statement (iii). We only consider the case when $b < |\beta_n|$ for all $n \in \mathcal{P}_+(N)$ since the proof of the other case is similar. We first conclude from (2.2) that $\{|A_n e^{i\alpha_n x_1 + i\beta_n (h-\sigma)}| : n \in \mathcal{P}_+(N)\}$ is uniformly bounded. Noting that in this case $(i\beta_n + b) < 0$ for all $n \in \mathcal{P}_+(N)$, we have

$$\begin{aligned} \sum_{n \in \mathcal{P}_+(N)} |A_n e^{i\alpha_n x_1 + i\beta_n x_2 + b x_2}| &\leq \sum_{n \in \mathcal{P}_+(N)} \left| A_n e^{i\alpha_n x_1 + i\beta_n (h-\sigma)} \right| e^{b(h-\sigma)} e^{(i\beta_n + b)(x_2 - h + \sigma)} \\ &\leq \sum_{n \in \mathcal{P}_+(N)} C e^{-|n|/C} \end{aligned}$$

uniformly for all $x \in U_h$, where we have used the fact that $\sigma \sqrt{(2n\pi/L + \alpha_0)^2 - k^2} - b > |n|/C$ holds for sufficiently large $|n|$ provided the constant $C > 0$ is large enough. This implies that (2.1) is uniformly and absolutely convergent for $x \in U_h$.

Finally, noting that

$$\left| \frac{1}{T} \int_T^{2T} a_j e^{ib_j t} dt \right| = \left| \frac{1}{T} \frac{a_j (e^{2ib_j T} - e^{ib_j T})}{ib_j} \right| \leq \frac{1}{T} \frac{2|a_j|}{|b_j|}, \quad j = 1, \dots, N,$$

it is easy to see that statement (iv) holds. □

LEMMA 2.2. Let $u(x; \theta_m)$ be the total field corresponding to the diffraction problem (1.1)–(1.4) with the incident angle $\theta = \theta_m \in (-\pi/2, \pi/2)$ for $m = 1, \dots, M$ and $M \in \mathbb{Z}_+$. Suppose $\{\theta_m\}_{m=1}^M$ are distinct incident angles. Then $\{u(x; \theta_m)\}_{m=1}^M$ are linearly independent in Ω .

Proof. Assume that $\sum_{m=1}^M c_m u(x; \theta_m) = 0$ in Ω for some $c_m \in \mathbb{C}$, $m = 1, \dots, M$. To indicate the dependence of u^s on the incident angle, we rewrite the Rayleigh expansion (1.4) as

$$u^s(x; \theta_m) = \sum_{n \in \mathbb{Z}} A_n(\theta_m) e^{i\alpha_n(\theta_m)x_1 + i\beta_n(\theta_m)x_2}, \quad x \in U_h,$$

where $h > \max\{x_2 : x \in \Gamma\}$ and $\alpha_n(\theta_m) := \alpha(\theta_m) + 2n\pi/L$ with $\alpha(\theta_m) := k \sin \theta_m$ and $\beta_n(\theta_m) \in \mathbb{C}$ are defined as in (1.5) with the incident angle $\theta = \theta_m$. Then, by (1.6) it follows that

$$(2.3) \quad \sum_{m=1}^M c_m u(x; \theta_m) = \sum_{m=1}^M c_m \left(\sum_{n \in \mathbb{Z} \cup \{i\}} A_n(\theta_m) e^{i\alpha_n(\theta_m)x_1 + i\beta_n(\theta_m)x_2} \right) = 0.$$

For any $\tilde{m} \in \{1, 2, \dots, M\}$, multiplying (2.3) by $e^{-i\beta_i(\theta_{\tilde{m}})x_2}$ we obtain

$$(2.4) \quad \sum_{m \in \mathcal{I}_{\tilde{m}}} c_m e^{i\alpha_i(\theta_m)x_1} + \sum_{m \in \{1, \dots, M\} \setminus \mathcal{I}_{\tilde{m}}} c_m e^{i\alpha_i(\theta_m)x_1 + i[\beta_i(\theta_m) - \beta_i(\theta_{\tilde{m}})]x_2} + \sum_{m=1}^M c_m \left(\sum_{n \in \mathbb{Z}} A_n(\theta_m) e^{i\alpha_n(\theta_m)x_1 + i[\beta_n(\theta_m) - \beta_i(\theta_{\tilde{m}})]x_2} \right) = 0, \quad x \in U_h,$$

where $\mathcal{I}_{\tilde{m}} := \{m \in \{1, \dots, M\} : \beta_i(\theta_m) = \beta_i(\theta_{\tilde{m}})\}$.

Next we claim that

$$(2.5) \quad \lim_{H \rightarrow +\infty} \frac{1}{H} \int_H^{2H} \sum_{m \in \{1, \dots, M\} \setminus \mathcal{I}_{\tilde{m}}} c_m e^{i\alpha_i(\theta_m)x_1 + i[\beta_i(\theta_m) - \beta_i(\theta_{\tilde{m}})]x_2} dx_2 = 0,$$

$$(2.6) \quad \lim_{H \rightarrow +\infty} \frac{1}{H} \int_H^{2H} \sum_{m=1}^M c_m \left(\sum_{n \in \mathbb{Z}} A_n(\theta_m) e^{i\alpha_n(\theta_m)x_1 + i[\beta_n(\theta_m) - \beta_i(\theta_{\tilde{m}})]x_2} \right) dx_2 = 0$$

for all $x_1 \in \mathbb{R}$. In fact, (2.5) follows easily from Lemma 2.1 (iv). To prove (2.6), let $m \in \{1, \dots, M\}$ be arbitrarily fixed. For $N > 0$ large enough we set $\mathcal{J}_1(N) := \{n \in \mathbb{Z} : |\alpha_n(\theta_m)| > k, |n| > N\}$ and $\mathcal{J}_2(N) := \{n \in \mathbb{Z} : |\alpha_n(\theta_m)| > k, |n| \leq N\}$. Using $|e^{-i\beta_i(\theta_{\tilde{m}})x_2}| = 1$, it follows from Lemma 2.1 (ii) that

$$(2.7) \quad \lim_{N \rightarrow +\infty} \sum_{n \in \mathcal{J}_1(N)} \left| A_n(\theta_m) e^{i\alpha_n(\theta_m)x_1 + i[\beta_n(\theta_m) - \beta_i(\theta_{\tilde{m}})]x_2} \right| = 0$$

uniformly for all $x \in U_h$. For any fixed $N \in \mathbb{Z}_+$, since $\mathcal{J}_2(N)$ is a finite set and $i\beta_n(\theta_m) < 0$ for all $n \in \mathcal{J}_2(N)$, we have

$$(2.8) \quad \lim_{x_2 \rightarrow +\infty} \sum_{n \in \mathcal{J}_2(N)} \left| A_n(\theta_m) e^{i\alpha_n(\theta_m)x_1 + i[\beta_n(\theta_m) - \beta_i(\theta_{\tilde{m}})]x_2} \right| = 0$$

uniformly for all $x_1 \in \mathbb{R}$. Since $\mathcal{J}_3 := \{n \in \mathbb{Z} : |\alpha_n(\theta_m)| \leq k\}$ is also a finite set and $\beta_n(\theta_m) \geq 0 > \beta_i(\theta_{\tilde{m}})$ for all $n \in \mathcal{J}_3$, it follows from Lemma 2.1 (iv) that

$$\lim_{H \rightarrow +\infty} \frac{1}{H} \int_H^{2H} \sum_{n \in \mathcal{J}_3} A_n(\theta_m) e^{i\alpha_n(\theta_m)x_1 + i[\beta_n(\theta_m) - \beta_i(\theta_{\tilde{m}})]x_2} dx_2 = 0$$

uniformly for all $x_1 \in \mathbb{R}$. This, together with (2.7)–(2.8), implies that (2.6) holds.

Combining (2.4)–(2.6), we arrive at

$$(2.9) \quad \sum_{m \in \mathcal{I}_{\tilde{m}}} c_m e^{i\alpha_i(\theta_m)x_1} = 0, \quad x_1 \in \mathbb{R}.$$

Multiplying (2.9) by $e^{-i\alpha_i(\theta_{\tilde{m}})x_1}$ we obtain

$$\sum_{m \in \mathcal{K}_{\tilde{m}}} c_m + \sum_{m \in \mathcal{I}_{\tilde{m}} \setminus \mathcal{K}_{\tilde{m}}} c_m e^{i[\alpha_i(\theta_m) - \alpha_i(\theta_{\tilde{m}})]x_1} = 0, \quad x_1 \in \mathbb{R},$$

where $\mathcal{K}_{\tilde{m}} := \{m \in \mathcal{I}_{\tilde{m}} : \alpha_i(\theta_m) = \alpha_i(\theta_{\tilde{m}})\}$. Obviously, $\mathcal{K}_{\tilde{m}} = \{\tilde{m}\}$. Then it follows from Lemma 2.1 (iv) that $c_{\tilde{m}} = 0$. By the arbitrariness of \tilde{m} it follows that $c_m = 0$ for all $m = 1, \dots, M$, implying that $\{u(x; \theta_m)\}_{m=1}^M$ are linearly independent in Ω . \square

Remark 2.1. By (1.6) the total field u to the diffraction problem (1.1)–(1.4) is given by

$$(2.10) \quad u(x) = \sum_{n \in \mathbb{Z} \cup \{i\}} A_n e^{i\alpha_n x_1 + i\beta_n x_2}, \quad x \in U_h.$$

We claim that

$$(2.11) \quad u \not\equiv 0 \text{ in } \Omega.$$

Assume to the contrary that $u \equiv 0$ in Ω . Then, proceeding as in the proof of Lemma 2.2, we first multiply (2.10) by $e^{-i\beta_i x_2}$ and then by $e^{-i\alpha_i x_1}$ to obtain that $A_i = 0$, which contradicts $A_i = 1$. This implies that (2.11) holds.

In the remaining part of this paper, we consider two periodic curves $\Gamma^{(1)}$ and $\Gamma^{(2)}$ with periods $L_1 > 0$ and $L_2 > 0$, respectively. Denote by Ω_j the unbounded connected domain above $\Gamma^{(j)}$ for $j = 1, 2$. Set $\Gamma_h := \{x : x_2 = h\}$ for some $h > \max\{x_2 : x \in \Gamma^{(1)} \cup \Gamma^{(2)}\}$. Denote by $u_j^s(x; \theta)$ and $u_j(x; \theta)$ the scattered field and total field, respectively, for incident plane wave $u^i(x; \theta)$ with $\theta \in (-\pi/2, \pi/2)$ corresponding to the curve $\Gamma^{(j)}$, $j = 1, 2$. Analogously, denote by $(\alpha_n^{(j)}, \beta_n^{(j)})$ the pair (α_n, β_n) (see (1.5) and (1.6)) and by $A_n^{(j)}$ the Rayleigh coefficient A_n in (1.4) and (1.6) corresponding to $\Gamma = \Gamma^{(j)}$ for $n \in \mathbb{Z} \cup \{i\}$ and $j = 1, 2$.

3. Determination of grating period from phased data. In this section we consider the inverse problem, that is, whether it is possible to determine the period of a periodic curve from phased near-field data corresponding to one incident plane wave. Since the total field u to the forward diffraction model (1.1)–(1.4) is required to be α -quasi-periodic, it is seen that $e^{-i\alpha x_1} u(x)$ is L -periodic with respect to x_1 . Actually, this is also implied by (1.4) and (1.6). However, the period L may not be the minimum period of $e^{-i\alpha x_1} u(x)$, as illustrated in the following remark, which presents two diffraction grating curves with different minimum periods which can generate identical near-field data for one incident plane wave. Such an example was motivated by the classification of unidentifiable polygonal diffraction gratings using one incident plane wave; see [8, 9, 16, 17].

Remark 3.1. Consider the example with $u = u^i + u^s$, where

$$(3.1) \quad u^i(x) = e^{i(-x_1 - \sqrt{3}x_2)}, \quad u^s(x) = e^{i(x_1 + \sqrt{3}x_2)} - e^{-2ix_1} - e^{2ix_1}.$$




FIG. 2. Contour of the total field u given by (3.2). The red solid line and the red dash-dot line denote two grating curves with different minimum periods, respectively.

Obviously, u^i is a plane wave defined as in (1.1) with incident angle $\theta = -\pi/6$ and wave number $k = 2$, implying that $\alpha = -1$. Note that if we choose the period $L = 2\pi$, then the Rayleigh frequency occurs (since $\beta_{-1} = \beta_3 = 0$ in this case). A straightforward calculation shows

$$(3.2) \quad u(x) = 2 \cos(x_1 + \sqrt{3}x_2) - 2 \cos(2x_1) = -4 \sin \frac{3x_1 + \sqrt{3}x_2}{2} \sin \frac{-x_1 + \sqrt{3}x_2}{2}.$$

Therefore, the zeros of $u(x)$ consist of two families of parallel lines,

$$\begin{aligned} l_n^{(1)} &:= \{x = (x_1, x_2) \in \mathbb{R}^2 : 3x_1 + \sqrt{3}x_2 = 2n\pi\}, \\ l_n^{(2)} &:= \{x = (x_1, x_2) \in \mathbb{R}^2 : -x_1 + \sqrt{3}x_2 = 2n\pi\} \end{aligned}$$

for $n \in \mathbb{Z}$, which form a grid in \mathbb{R}^2 , as illustrated by Figure 2. It is obvious that the two curves $\Gamma^{(1)}$ and $\Gamma^{(2)}$ plotted by the red solid line ‘-’ and the red dashed line ‘-.’, respectively, as shown in Figure 2, lie on the above grid. The minimum period of $\Gamma^{(1)}$ and $\Gamma^{(2)}$ is $L_1 = 2\pi$ and $L_2 = 4\pi$, respectively. From the above discussions and the formula (3.1), it can be seen that u^s is the scattered field to the diffraction problem (1.1)–(1.4), with the curve $\Gamma = \Gamma^{(1)}$ and the period $L = L_1$, and satisfies the Rayleigh expansion (1.4) with nonzero Rayleigh coefficients $A_2^{(1)} = 1$, $A_{-1}^{(1)} = A_3^{(1)} = -1$. However, on the other hand, it is also easily seen that u^s is the scattered field to the diffraction problem (1.1)–(1.4), with the curve $\Gamma = \Gamma^{(2)}$ and the period $L = L_2$, and satisfies the Rayleigh expansion (1.4) with nonzero Rayleigh coefficients $A_4^{(2)} = 1$, $A_{-2}^{(2)} = A_6^{(2)} = -1$. This example shows that it is impossible to determine the minimum period (also the shape) of a grating curve from phased near-field data corresponding to one incident plane wave.

In general, one can only find a common period of two grating curves if their scattered fields coincide. This will be proved rigorously in Theorem 3.1, where the periodic curves do not need to satisfy the smoothness Conditions (i) and (ii).

THEOREM 3.1. *Suppose $\theta \in (-\pi/2, \pi/2)$ is an arbitrarily fixed incident angle. Let $\Gamma^{(1)}$ and $\Gamma^{(2)}$ be two periodic curves. If the corresponding scattered fields satisfy*

$$(3.3) \quad u_1^s(x; \theta) = u_2^s(x; \theta) \quad \text{on} \quad x_2 = h > \max\{x_2 : x \in \Gamma^{(1)} \cup \Gamma^{(2)}\},$$

then there exists $L > 0$ such that L is a period of both $\Gamma^{(1)}$ and $\Gamma^{(2)}$.

Proof. Suppose $L_j > 0$ is a period of the curve $\Gamma^{(j)}$, $j = 1, 2$. Then the corresponding scattered field $u_j^s(x; \theta)$ satisfies the Rayleigh expansions

$$(3.4) \quad u_j^s(x) = \sum_{n \in \mathbb{Z}} A_n^{(j)} e^{i\alpha_n^{(j)} x_1 + i\beta_n^{(j)} x_2}, \quad x \in U_h := \{x \in \mathbb{R}^2 : x_2 > h\}, j = 1, 2,$$

where $\alpha_n^{(j)}, \beta_n^{(j)}$, and the coefficients $A_n^{(j)}$ that depend on k, θ , and $\Gamma^{(j)}$ are defined analogously to α_n, β_n , and A_n , with Γ replaced by $\Gamma^{(j)}$. Note that the following conditions are fulfilled:

- (i) $u_1^s - u_2^s$ satisfies the Helmholtz equation in U_h ;
- (ii) $u_1^s - u_2^s = 0$ on $\Gamma_h := \{x : x_2 = h\}$;
- (iii) $\sup_{x \in U_h} |u_1^s(x) - u_2^s(x)| < +\infty$;
- (iv) $u_1^s - u_2^s$ satisfies the upward propagating radiation condition (see [13, Definition 2.2]).

In fact, (i) follows from (1.1) and (1.2), and (ii) follows from (3.3). (iii) and (iv) are implied by the Rayleigh expansions (3.4) (see Lemma 2.1 (i) and [13, p. 1777]). By uniqueness to the Dirichlet boundary value problem in U_h (see [13, Theorem 3.4]), it follows that

$$(3.5) \quad u_1^s(x; \theta) = u_2^s(x; \theta), \quad x \in U_h.$$

We now consider the following two cases.

Case 1. L_1/L_2 is rational.

Let $p/q = L_1/L_2$ with reduced fraction p/q and positive integers $p, q \in \mathbb{Z}_+$. Set $L := qL_1$. Then $L = pL_2$. Thus L is a common period for both $\Gamma^{(1)}$ and $\Gamma^{(2)}$.

Case 2. L_1/L_2 is irrational.

We claim that any $L > 0$ is a period of both $\Gamma^{(1)}$ and $\Gamma^{(2)}$. To do this, we first deduce from the fact that L_1/L_2 is irrational that

$$(3.6) \quad \alpha_m^{(1)} \neq \alpha_n^{(2)} \text{ for all } (m, n) \in \mathbb{Z}^2 \setminus \{(0, 0)\} \text{ and } \alpha_0^{(1)} = \alpha_0^{(2)} = k \sin \theta.$$

It follows from (3.4) and (3.5) that

$$(3.7) \quad \sum_{n \in \mathbb{Z}} A_n^{(1)} e^{i\alpha_n^{(1)}x_1 + i\beta_n^{(1)}x_2} - \sum_{n \in \mathbb{Z}} A_n^{(2)} e^{i\alpha_n^{(2)}x_1 + i\beta_n^{(2)}x_2} = 0, \quad x \in U_h.$$

The proof of this case can be divided into three steps as follows.

Step 1. We prove that

$$(3.8) \quad A_0^{(1)} = A_0^{(2)},$$

$$(3.9) \quad A_n^{(1)} = 0 \text{ for all } n \in \mathbb{Z} \setminus \{0\} \text{ such that } |\alpha_n^{(1)}| \leq k,$$

$$(3.10) \quad A_n^{(2)} = 0 \text{ for all } n \in \mathbb{Z} \setminus \{0\} \text{ such that } |\alpha_n^{(2)}| \leq k.$$

Let $\tilde{n} \in \mathbb{Z}$ be arbitrarily fixed such that $|\alpha_{\tilde{n}}^{(1)}| \leq k$. Multiplying (3.7) by $e^{-i\beta_{\tilde{n}}^{(1)}x_2}$, we obtain

$$(3.11) \quad \sum_{n \in \mathcal{I}_{\tilde{n}}^{(1)}} A_n^{(1)} e^{i\alpha_n^{(1)}x_1} + \sum_{n \in \mathbb{Z} \setminus \mathcal{I}_{\tilde{n}}^{(1)}} A_n^{(1)} e^{i\alpha_n^{(1)}x_1 + i(\beta_n^{(1)} - \beta_{\tilde{n}}^{(1)})x_2} - \sum_{n \in \mathcal{I}_{\tilde{n}}^{(2)}} A_n^{(2)} e^{i\alpha_n^{(2)}x_1} - \sum_{n \in \mathbb{Z} \setminus \mathcal{I}_{\tilde{n}}^{(2)}} A_n^{(2)} e^{i\alpha_n^{(2)}x_1 + i(\beta_n^{(2)} - \beta_{\tilde{n}}^{(1)})x_2} = 0, \quad x \in U_h,$$

where $\mathcal{I}_{\tilde{n}}^{(j)} := \{n \in \mathbb{Z} : \beta_n^{(j)} = \beta_{\tilde{n}}^{(1)}\}$ is at most a finite set for $j = 1, 2$. Analogously to (2.6), using $|e^{i\beta_{\tilde{n}}^{(1)}x_2}| = 1$, we can apply Lemma 2.1 (ii) and (iv) to obtain

$$\lim_{H \rightarrow +\infty} \frac{1}{H} \int_H^{2H} \sum_{n \in \mathbb{Z} \setminus \mathcal{I}_{\tilde{n}}^{(j)}} A_n^{(j)} e^{i\alpha_n^{(j)}x_1 + i(\beta_n^{(j)} - \beta_{\tilde{n}}^{(1)})x_2} dx_2 = 0, \quad j = 1, 2,$$

for all $x_1 \in \mathbb{R}$. Therefore, it follows from (3.11) that

$$(3.12) \quad \sum_{n \in \mathcal{I}_{\tilde{n}}^{(1)}} A_n^{(1)} e^{i\alpha_n^{(1)} x_1} - \sum_{n \in \mathcal{I}_{\tilde{n}}^{(2)}} A_n^{(2)} e^{i\alpha_n^{(2)} x_1} = 0, \quad x_1 \in \mathbb{R}.$$

Similarly, multiplying (3.12) by $e^{-i\alpha_{\tilde{n}}^{(1)} x_1}$, we can deduce from Lemma 2.1 (iv) that

$$(3.13) \quad \sum_{n \in \mathcal{K}_{\tilde{n}}^{(1)}} A_n^{(1)} - \sum_{n \in \mathcal{K}_{\tilde{n}}^{(2)}} A_n^{(2)} = 0,$$

where $\mathcal{K}_{\tilde{n}}^{(j)} := \{n \in \mathbb{Z} : \alpha_n^{(j)} = \alpha_{\tilde{n}}^{(j)}, \beta_n^{(j)} = \beta_{\tilde{n}}^{(j)}\}$, $j = 1, 2$. Obviously, $\mathcal{K}_{\tilde{n}}^{(1)} = \{\tilde{n}\}$. In view of (3.6), we know that $\mathcal{K}_{\tilde{n}}^{(2)} = \{0\}$ if $\tilde{n} = 0$ and $\mathcal{K}_{\tilde{n}}^{(2)} = \emptyset$ if $\tilde{n} \in \mathbb{Z} \setminus \{0\}$. These, together with (3.13), imply (3.8) and (3.9). By interchanging the roles of u_1^s and u_2^s , we can employ an argument similar to that above to obtain (3.10).

Step 2. We prove that

$$(3.14) \quad A_n^{(1)} = 0 \text{ for all } n \in \mathbb{Z} \text{ such that } |\alpha_n^{(1)}| > k,$$

$$(3.15) \quad A_n^{(2)} = 0 \text{ for all } n \in \mathbb{Z} \text{ such that } |\alpha_n^{(2)}| > k.$$

Set $\mathcal{P}^{(j)} := \{n \in \mathbb{Z} : |\alpha_n^{(j)}| > k\}$, $j = 1, 2$. It follows from (3.7)–(3.10) that

$$(3.16) \quad \sum_{n \in \mathcal{P}^{(1)}} A_n^{(1)} e^{i\alpha_n^{(1)} x_1 + i\beta_n^{(1)} x_2} - \sum_{n \in \mathcal{P}^{(2)}} A_n^{(2)} e^{i\alpha_n^{(2)} x_1 + i\beta_n^{(2)} x_2} = 0, \quad x \in U_h.$$

By (1.5), we can rearrange the elements in $\{(1, n) : n \in \mathcal{P}^{(1)}\} \cup \{(2, n) : n \in \mathcal{P}^{(2)}\}$ as a sequence $\{(p_\ell, q_\ell)\}_{\ell \in \mathbb{Z}_+}$ such that $\beta_{q_\ell}^{(p_\ell)} = i b_\ell$ with $b_\ell > 0$ and $b_\ell \leq b_{\ell+1}$ for all $\ell \in \mathbb{Z}_+$. Obviously, $b_\ell \rightarrow +\infty$ as $\ell \rightarrow +\infty$.

Without loss of generality, we may assume that $p_1 = 1$ and $q_1 = \tilde{n}$ for some $\tilde{n} \in \mathcal{P}^{(1)}$ and thus that $\beta_{q_1}^{(p_1)} = \beta_{\tilde{n}}^{(1)}$. Let $\mathcal{I}_{\tilde{n}}^{(j)}$ ($j = 1, 2$) be defined as in Step 1. It is clear that $\mathcal{I}_{\tilde{n}}^{(j)} = \{n \in \mathcal{P}^{(j)} : \beta_n^{(j)} = \beta_{\tilde{n}}^{(j)}\}$ and is at most a finite set. Then, multiplying (3.16) by $e^{-i\beta_{\tilde{n}}^{(1)} x_2}$, we obtain

$$(3.17) \quad \begin{aligned} & \sum_{n \in \mathcal{I}_{\tilde{n}}^{(1)}} A_n^{(1)} e^{i\alpha_n^{(1)} x_1} + \sum_{n \in \mathcal{P}^{(1)} \setminus \mathcal{I}_{\tilde{n}}^{(1)}} A_n^{(1)} e^{i\alpha_n^{(1)} x_1 + i(\beta_n^{(1)} - \beta_{\tilde{n}}^{(1)}) x_2} \\ & - \sum_{n \in \mathcal{I}_{\tilde{n}}^{(2)}} A_n^{(2)} e^{i\alpha_n^{(2)} x_1} - \sum_{n \in \mathcal{P}^{(2)} \setminus \mathcal{I}_{\tilde{n}}^{(2)}} A_n^{(2)} e^{i\alpha_n^{(2)} x_1 + i(\beta_n^{(2)} - \beta_{\tilde{n}}^{(1)}) x_2} = 0, \quad x \in U_h. \end{aligned}$$

For $N > 0$ large enough and $j = 1, 2$, we set $\mathcal{Q}_1^{(j)}(N) := \{n \in \mathcal{P}^{(j)} \setminus \mathcal{I}_{\tilde{n}}^{(j)} : |n| > N\}$ and $\mathcal{Q}_2^{(j)}(N) := \{n \in \mathcal{P}^{(j)} \setminus \mathcal{I}_{\tilde{n}}^{(j)} : |n| \leq N\}$. By Lemma 2.1 (iii), we have

$$(3.18) \quad \lim_{N \rightarrow +\infty} \sum_{n \in \mathcal{Q}_1^{(j)}(N)} \left| A_n^{(j)} e^{i\alpha_n^{(j)} x_1 + i(\beta_n^{(j)} - \beta_{\tilde{n}}^{(1)}) x_2} \right| = 0, \quad j = 1, 2,$$

uniformly for all $x \in U_h$. For any fixed $N > 0$, since $\mathcal{Q}_2^{(j)}(N)$ is a finite set and $i(\beta_n^{(j)} - \beta_{\tilde{n}}^{(1)}) < 0$ for all $n \in \mathcal{Q}_2^{(j)}(N)$ due to the definition of $\beta_{\tilde{n}}^{(1)}$, thus we have

$$(3.19) \quad \lim_{x_2 \rightarrow +\infty} \sum_{n \in \mathcal{Q}_2^{(j)}(N)} \left| A_n^{(j)} e^{i\alpha_n^{(j)} x_1 + i(\beta_n^{(j)} - \beta_{\tilde{n}}^{(1)}) x_2} \right| = 0, \quad j = 1, 2,$$

uniformly for all $x_1 \in \mathbb{R}$. Thus, it follows from (3.18) and (3.19) that

$$\lim_{x_2 \rightarrow +\infty} \sum_{n \in \mathcal{P}^{(j)} \setminus \mathcal{I}_n^{(j)}} A_n^{(j)} e^{i\alpha_n^{(j)} x_1 + i(\beta_n^{(j)} - \beta_n^{(1)}) x_2} = 0, \quad j = 1, 2,$$

for all $x_1 \in \mathbb{R}$. This, together with (3.17), implies that (3.12) holds. Analogously to Step 1, multiplying (3.12) by $e^{-i\alpha_n^{(1)} x_1}$, we can apply Lemma 2.1 (iv) to obtain (3.13) and thus $A_n^{(1)} = A_{q_1}^{(p_1)} = 0$. Taking this into (3.16), we obtain that (3.16) holds with $\mathcal{P}^{(1)}$ replaced by $\mathcal{P}^{(1)} \setminus \{q_1\}$. Then using the same argument as above, we can obtain that $A_{q_2}^{(p_2)} = 0$. Now, we can repeat the same argument again to obtain that $A_{q_\ell}^{(p_\ell)} = 0$ for all $\ell \in \mathbb{Z}_+$. This means that (3.14) and (3.15) hold.

Step 3. Combining (3.8)–(3.10), (3.14), and (3.15), we arrive at

$$A_0^{(1)} = A_0^{(2)} \text{ and } A_n^{(1)} = A_n^{(2)} = 0 \text{ for } n \in \mathbb{Z} \setminus \{0\}.$$

Then by the Dirichlet boundary condition imposed on $\Gamma^{(j)}$ ($j = 1, 2$), we have

$$e^{i\alpha_0 x_1 - i\beta_0 x_2} = u^i(x) = -u_j^s(x) = -A_0^{(j)} e^{i\alpha_0 x_1 + i\beta_0 x_2}, \quad x \in \Gamma^{(j)}, \quad j = 1, 2.$$

This further implies that $\Gamma^{(j)}$ ($j = 1, 2$) is a straight line parallel to the x_1 -axis since $A_0^{(j)}$ is a constant. Thus, any $L > 0$ is a common period of $\Gamma^{(1)}$ and $\Gamma^{(2)}$. \square

Remark 3.2. Case (ii) in the proof of Theorem 3.1 implies that if the relation (3.3) holds and one of $\Gamma^{(j)}$ ($j = 1, 2$) is not a straight line parallel to the x_1 -axis, then it is impossible that L_1/L_2 is irrational.

4. Uniqueness with phased data. In this section, we prove that a periodic curve with Dirichlet boundary condition fulfilling Condition (i) or Condition (ii) can be uniquely determined by the fixed-frequency near-field data corresponding to incident plane waves with distinct angles (i.e., Theorem 1.1 with phased data). This differs from [23], where fixed-direction incident plane waves with different frequencies are used, and this also differs from [29] which involves fixed-frequency quasi-periodic incident waves with the same phase shift. For the inverse problem to recovering a periodic curve from near-field data corresponding to incident plane waves with distinct directions, difficulties arise from the fact that the corresponding total fields have different phase shifts since $\alpha = k \sin \theta$ depends on the incident angle θ . We rephrase Theorem 1.1 with phased data in Theorem 4.1 below, which is the main uniqueness result of this section. Here we shall provide a proof based on the ideas of Schiffer both for bounded obstacles (see [15]) and for periodic structures with multifrequency data (see [23]) and the concept of dispersion relations (see, e.g., [22, 36, 40]) arising from the analysis of photonic crystals.

THEOREM 4.1. *Let $\Gamma^{(1)}$ and $\Gamma^{(2)}$ be two periodic curves with Dirichlet boundary conditions. Assume both satisfy Condition (i) or both satisfy Condition (ii). Suppose that the periods of $\Gamma^{(1)}$ and $\Gamma^{(2)}$ are unknown. If the corresponding total fields satisfy*

$$(4.1) \quad u_1(x; \theta_n) = u_2(x; \theta_n), \quad x \in \mathcal{S}, \quad n \in \mathbb{Z}_+,$$

where $\{\theta_n\}_{n=1}^\infty$ are distinct incident angles in $(-\pi/2, \pi/2)$, then $\Gamma^{(1)} = \Gamma^{(2)}$. Here, $\mathcal{S} \subset \Gamma_h$ is a line segment with $\Gamma_h := \{x : x_2 = h\}$ and with $h > \max\{x_2 : x \in \Gamma^{(1)} \cup \Gamma^{(2)}\}$ being an arbitrary constant.

Downloaded 03/18/23 to 202.113.19.215 . Redistribution subject to SIAM license or copyright; see https://pubs.siam.org/terms-privacy




FIG. 3. The bounded domain D in Case (i): $\Gamma^{(1)} \cap \Gamma^{(2)} \neq \emptyset$.

Remark 4.1. In Theorem 4.1, the wave number $k > 0$ is fixed, and $u_j(\cdot; \theta_n)$ ($j = 1, 2$) are forward solutions corresponding to $\Gamma^{(j)}$ and the incident direction θ_n , respectively. The uniqueness result of Theorem 4.1 remains true even if the incident angle causes a Rayleigh frequency, i.e., $k^2 = (k \sin \theta_n + m)^2$ for some $n \in \mathbb{Z}_+$ and $m \in \mathbb{Z}$. The preliminary results presented in sections 2 and 3 and the proof of Theorem 4.1 are all valid in the case of a Rayleigh frequency.

Since u_1 and u_2 are analytic functions of $x \in \Gamma_h$, (4.1) is equivalent to $u_1(x; \theta_n) = u_2(x; \theta_n)$ for all $x \in \Gamma_h$ and $n \in \mathbb{Z}_+$. Therefore, $u_1^s(x; \theta_n) = u_2^s(x; \theta_n)$ for all $x \in \Gamma_h$ and $n \in \mathbb{Z}_+$. Analogously to (3.5), we have $u_1^s(x; \theta_n) = u_2^s(x; \theta_n)$ for all $x \in U_h$ and $n \in \mathbb{Z}_+$. By analyticity we arrive at

$$(4.2) \quad u_1^s(x; \theta_n) = u_2^s(x; \theta_n), \quad x \in \Omega', n \in \mathbb{Z}_+,$$

where Ω' denotes the unbounded component of $\Omega_1 \cap \Omega_2$ which can be connected to U_h . By Theorem 3.1, the above relation (4.2) implies that there exists $L > 0$ such that L is a common period of $\Gamma^{(1)}$ and $\Gamma^{(2)}$. Without loss of generality, we may assume $L = 2\pi$ in the rest of this section. Assume to the contrary that $\Gamma^{(1)} \neq \Gamma^{(2)}$. We need to consider the following two cases:

$$\text{Case (i): } \Gamma^{(1)} \cap \Gamma^{(2)} \neq \emptyset; \quad \text{Case (ii): } \Gamma^{(1)} \cap \Gamma^{(2)} = \emptyset.$$

The proofs of Theorem 4.1 for these two cases will be given in the following subsections.

4.1. Proof of Theorem 4.1 for Case (i): $\Gamma^{(1)} \cap \Gamma^{(2)} \neq \emptyset$. Since $\Gamma^{(1)} \cap \Gamma^{(2)} \neq \emptyset$ and both $\Gamma^{(1)}$ and $\Gamma^{(2)}$ are 2π -periodic, there exists at least one bounded domain D enclosed by $\Gamma^{(1)}$ and $\Gamma^{(2)}$. In other words, $\partial D \subset \Gamma^{(1)} \cup \Gamma^{(2)}$. Without loss of generality, we may suppose that $D \subset \Omega_1 \setminus \bar{\Omega}'$ as shown in Figure 3. It follows from Remark 2.1, formula (4.2), and the Dirichlet boundary condition of $u_j(x; \theta_n)$ on $\Gamma^{(j)}$ that the total field $u_1(x; \theta_n) := u^i(x; \theta_n) + u_1^s(x; \theta_n)$ is a nontrivial solution to the eigenvalue problem

$$\Delta u + k^2 u = 0 \quad \text{in } D, \quad u = 0 \quad \text{on } \partial D$$

for all $n \in \mathbb{Z}_+$. In other words, $u_1(x; \theta_n)$ is a Dirichlet eigenfunction of the negative Laplacian in D for each $n \in \mathbb{Z}_+$. Recall from Lemma 2.2 that $\{u_1(x; \theta_n)\}_{n=1}^N$ are linearly independent functions in D for any positive integer $N < +\infty$. However, by a similar argument as in the proof of [14, Theorem 5.1], it follows that there are at most finitely many independent Dirichlet eigenfunctions of the negative Laplacian in $H_0^1(D)$ corresponding to the eigenvalue $k^2 > 0$. This contradiction implies that Case (i) does not hold.

Remark 4.2. It should be noted that the proof of [14, Theorem 5.1] relies essentially on the a priori estimate of solutions after the Gram-Schmidt orthogonalization of $\{u_1(x; \theta_n)\}_{n \in \mathbb{Z}_+}$ (see [14, third formula on page 140]). However, if D were an unbounded periodic strip, as will be seen in Case (ii), it would be difficult to establish




FIG. 4. The unbounded periodic strip D and its one periodicity cell M in Case (ii): $\Gamma^{(1)} \cap \Gamma^{(2)} = \emptyset$.

an analogous a priori estimate of solutions with different incident angles (or, equivalently, with different phase shifts $k \sin \theta_n$) after the Gram–Schmidt orthogonalization. Hence, the aforementioned arguments cannot be used for treating Case (ii).

4.2. Proof of Theorem 4.1 for Case (ii): $\Gamma^{(1)} \cap \Gamma^{(2)} = \emptyset$. We suppose without loss of generality that $\Gamma^{(2)}$ lies entirely above $\Gamma^{(1)}$ as shown in Figure 4. Denote by D the unbounded 2π -periodic strip (waveguide) lying between the two curves. To investigate the dependence of solutions on the quasi-periodic shift $\alpha = \alpha(\theta_n) = k \sin \theta_n$, we set $w_n(x) := e^{-i\alpha(\theta_n)x_1} u_1(x; \theta_n)$. It then follows from (1.2) and (4.2) that w_n satisfies the periodic boundary value problem

$$(4.3) \quad \begin{cases} \nabla_{\alpha(\theta_n)} \cdot \nabla_{\alpha(\theta_n)} w_n + k^2 w_n = 0 & \text{in } D, \\ w_n = 0 & \text{on } \Gamma^{(1)} \cup \Gamma^{(2)}, \\ w_n \text{ is } 2\pi\text{-periodic with respect to } x_1 & \text{in } D, \end{cases}$$

for all $n \in \mathbb{Z}_+$, where $\nabla_{\alpha(\theta_n)} := (\partial_1 + i\alpha(\theta_n), \partial_2)^\top$. For $\alpha = k\mu$ with $\mu \in (-1, 1)$, we consider the following abstract Dirichlet boundary value problem in a closed periodic waveguide D :

$$(BVP) \quad \begin{cases} \nabla_\alpha \cdot \nabla_\alpha w + k^2 w = 0 & \text{in } D, \\ w = 0 & \text{on } \Gamma^{(1)} \cup \Gamma^{(2)}, \\ w \text{ is } 2\pi\text{-periodic with respect to } x_1 & \text{in } D. \end{cases}$$

DEFINITION 4.1. For any fixed $k > 0$, we say that $\mu \in (-1, 1)$ is called a μ -eigenvalue if (BVP) admits a nontrivial solution in the space $H_{0,0}^1(D) := \{w \in H_{loc}^1(D) : w \text{ is } 2\pi\text{-periodic with respect to } x_1, w = 0 \text{ on } \partial D\}$. Accordingly, the nontrivial solution is the associated eigenfunction.

Since $u_1(x; \theta_n) \not\equiv 0$ for $x \in \Omega_1$ (see Remark 2.1), we conclude from (4.3) that $\sin \theta_n$ is a μ -eigenvalue to (BVP) with the eigenfunction w_n for all $n \in \mathbb{Z}_+$. On the other hand, for any fixed $\mu \in (-1, 1)$, we say that $k > 0$ is called a k -eigenvalue if (BVP) admits a nontrivial solution $w \in H_{0,0}^1(D)$. As shown in [23, Theorem 2.3], the k -eigenvalues form a discrete set on the positive real-axis with the only accumulating point at infinity, and the associated eigenspace for each k -eigenvalue is of finite dimension. It is easy to observe that if $w(x)$ solves (BVP) with $\mu \in (-1, 1)$ and some $k_j(\mu)$, then the conjugate \bar{w} is also a nontrivial solution corresponding to $-\mu$. This implies the even symmetry of $k_j(\mu)$ with respect to the line $\mu = 0$, that is, $k_j(\mu) = k_j(-\mu)$ for each $\mu \in (-1, 1)$.

The α -dependent partial differential equation in (BVP) can be regarded as the Floquet–Bloch (FB) transform of the Helmholtz equation $(\Delta + k^2)u = 0$ in the x_1 -direction with the variable $\alpha \in \mathbb{R}$; see [22, 36]. The Bloch theory in one direction was well summarized in [22, section 3] for deriving physically meaningful radiation conditions in a closed periodic waveguide.

Let us now recall the dispersion relations for the 2π -periodic system (BVP), where the FB transform variable $\alpha \in \mathbb{R}$ is independent of k . For each $\alpha \in \mathbb{R}$, there also exists a discrete set of numbers $K_j(\alpha) > 0$ such that (BVP) admits nontrivial solutions with $k^2 = K_j(\alpha)$ for each $j = 1, 2, \dots$ (see Remark 4.3). By [27, Chapter 7], the function $\alpha \rightarrow K_j(\alpha)$ is continuous and piecewise analytic. Further, $K_j(\alpha)$ is not analytic at $\alpha = \alpha_0$ only if $k^2 = K_j(\alpha_0)$ is not a simple eigenvalue. Recall from (1.3) with $L = 2\pi$ that an α -quasi-periodic function must also be $(\alpha + j)$ -quasi-periodic for any $j \in \mathbb{N}$. It is easy to conclude that $K_j(\alpha) : \mathbb{R} \rightarrow \mathbb{R}$ is periodic in α with periodicity one. Restricting to one periodic interval $[-1/2, 1/2]$, we also have the even symmetry $K_j(\alpha) = K_j(-\alpha)$ for all $\alpha \in [-1/2, 1/2]$. The α -dependent eigenvalues $K_j(\alpha)$ can be relabeled for $j \in \mathbb{Z}_+$ so as to make the eigenvalues and associated eigenfunctions analytic in $\alpha \in \mathbb{R}$ (see, e.g., [27, Theorem 3.9, Chapter 7] or [22, section 3.3]). For $j \in \mathbb{Z}_+$ the curves given by $K_j(\alpha) : (-1/2, 1/2] \rightarrow \mathbb{R}$ for the relabeled indices are well known as dispersion relations, and the graphs of the dispersion relations define the Bloch variety [36]. Note that the dispersion curves are no longer periodic. Below we characterize the relation between the function $\mu \mapsto k(\mu)$ and the dispersion relation $\alpha \mapsto K(\alpha)$.

LEMMA 4.1. (i) *The function $k_j(\mu) : (-1, 1) \rightarrow \mathbb{R}_+$ must fulfill the dispersion relation $K_{j'}(\mu k_j(\mu)) = k_j^2(\mu)$ for some $j' \in \mathbb{Z}_+$. Conversely, from the dispersion relation $K_{j'}(\mu k) = k^2$ one can always deduce the function $k = k_j(\mu)$ for some $j \in \mathbb{Z}_+$.*

(ii) *If $k_j(\mu) \equiv \text{Const}$ for some $j \in \mathbb{Z}_+$, then $K_{j'}(\alpha) \equiv \text{Const}$ for some $j' \in \mathbb{Z}_+$ and vice versa.*

Proof. (i) The first part follows straightforwardly from the definitions of k_j and $K_{j'}$. To prove the second part, we set $F(k) := K(\mu k) - k^2$. Obviously, $dF/dk = \mu K'(\mu k) - 2k$, where $K'(\alpha) := dK/d\alpha$. If

$$(4.4) \quad K(\mu k) - k^2 = 0, \quad \mu K'(\mu k) - 2k = 0,$$

we can conclude that

$$\alpha K'(\alpha) - 2K(\alpha) = 0, \quad \alpha = \mu k.$$

Hence, $K(\alpha) = c\alpha^2$ for some constant $c \in \mathbb{R}$. By the 1-periodicity of K we obtain $c = 0$ and thus $K \equiv 0$. This further leads to $k = 0$, and by integration by part, any solution to (BVP) must vanish identically. Hence, the two relations in (4.4) cannot hold simultaneously. By the implicit function theorem one can always get the function $k = k_j(\mu)$ for some $j \in \mathbb{Z}_+$ from the dispersion relation $K_{j'}(\alpha) = k^2$.

(ii) The second assertion is a direct consequence of the first assertion. \square

Remark 4.3. We consider a special case when $D = \mathbb{R} \times (0, h)$ is a straight strip with some $h > 0$. By separation of variables, it was proved in [23] that the dispersion relation is given by

$$(4.5) \quad K_{n,m}(\alpha) = (\alpha + n)^2 + \left(\frac{m\pi}{h}\right)^2, \quad n \in \mathbb{Z}, m \in \mathbb{Z}_+,$$

when $|\alpha| < k$ (see [23, (3.5)]). By the same argument as in [23], (4.5) holds for all $\alpha \in \mathbb{R}$. Here, the dispersion relation $\{K_{n,m}(\alpha)\}_{n \in \mathbb{Z}, m \in \mathbb{Z}_+}$ is the rearrangement of $\{K_j(\alpha)\}_{j \in \mathbb{Z}_+}$ mentioned above.

For a proof of Theorem 4.1 in Case (ii), it suffices to prove that the μ -eigenvalues must be discrete for any fixed $k > 0$. To this end, we need the following proposition.

PROPOSITION 4.1. *Assume both $\Gamma^{(1)}$ and $\Gamma^{(2)}$ satisfy Condition (i) or both satisfy Condition (ii) such that $\Gamma^{(1)} \cap \Gamma^{(2)} = \emptyset$. Then the problem (BVP) has no flat dispersion curves, that is, $K_j(\alpha) \not\equiv \text{const}$ for any $j \in \mathbb{Z}_+$.*

The result of Proposition 4.1 was essentially contained in the proof of [40, Theorem 2.3] for general periodic partial differential equations in an open or closed waveguide. In a closed waveguide, both the Dirichlet and Neumann boundary conditions were considered there. Moreover, Proposition 4.1 applies to general 3-admissible periodic domains (see [40, Definition 2.2]), which can be obtained from a straight strip by a periodic $W^{3,\infty}$ -mapping/a 3-admissible mapping, including the periodic strips stated in Proposition 4.1. As a direct consequence of Proposition 4.1, we have the following result.

COROLLARY 4.1. *Let $k > 0$ be an arbitrarily fixed wave number. Under the conditions of Proposition 4.1, there exists at least one parameter $\mu \in (-1, 1)$ such that the periodic boundary value problem (BVP) admits the trivial solution only.*

Proof. Assume to the contrary that for some $k > 0$ the periodic boundary value problem (BVP) admits nontrivial solutions for each $\mu \in (-1, 1)$. This implies that $k_j(\mu) = k > 0$ for all $\mu \in (-1, 1)$ and for some $j \in \mathbb{Z}_+$. By Lemma 4.1 (ii), there exists one flat dispersion curve $K_{j'}(\alpha) \equiv k^2$ for some $j' \in \mathbb{Z}_+$ for the system (BVP), which contradicts Proposition 4.1. \square

In the case when $\alpha \in \mathbb{C}$ and $\text{Im } \alpha > 0$ are sufficiently large, the strict coercivity of the sesquilinear form corresponding to (BVP) was justified in the proof of [40, Theorem 3.4] contained in [40, section 5]. The proof was based on a suitable change of variables which reduces the α -eigenvalue problem over 3-admissible periodic domains to an equivalent problem over straight strips. This, together with the analytic Fredholm theory or the perturbation theory (see, e.g., [27, Chapter 7, Theorems 1.10 and 1.9] or [39, Theorem XIII.86]) and Lemma 4.1, also implies Corollary 4.1. Now, we state the discreteness of the μ -eigenvalues for any fixed $k > 0$ and complete the proof of Theorem 4.1 in Case (ii).

LEMMA 4.2. *Under the conditions of Proposition 4.1, the μ -eigenvalues of (BVP) form at most a discrete set in $(-1, 1)$ without any accumulating points on the real-axis.*

Proof. We carry out the proof following the ideas in the proof of [23, Theorem 2.3], where the k -eigenvalue problem was investigated for the case when μ is fixed. Let w be a solution to the problem (BVP). Let $M := \{x \in D : 0 < x_1 < 2\pi\}$ be one 2π -periodic cell (see Figure 4 for the geometry of M), and let H be the completion of $\{\varphi \in C_p^1(\overline{M}) : \varphi = 0 \text{ on } \partial D \cap \overline{M}\}$ with respect to the H^1 -norm, where C_p^1 denotes the space of differentiable functions which are 2π -periodic with respect to x_1 . Note that M may be disconnected. Then we can apply Green's theorem to obtain that for any function $\psi \in H$,

$$(4.6) \quad \int_M \nabla w \cdot \nabla \overline{\psi} dx + \mu \int_M (-2ik \partial_1 w \overline{\psi}) dx + (\mu^2 - 1) \int_M k^2 w \overline{\psi} dx = 0.$$

Let $\langle \cdot, \cdot \rangle_H$ denote the inner product of the Hilbert space H , which is given by

$$\langle \varphi, \psi \rangle_H := \int_M \nabla \varphi \cdot \nabla \overline{\psi} dx, \quad \varphi, \psi \in H.$$

By Poincaré's inequality, it is known that $\langle \cdot, \cdot \rangle_H$ is equivalent to the ordinary inner product in $H^1(M)$. Then with the aid of Riesz's representation theorem, there exist $B, C \in \mathcal{L}(H)$ such that

$$\int_M (-2ik\partial_1\varphi\bar{\psi}) dx = \langle B\varphi, \psi \rangle_H, \quad \varphi, \psi \in H,$$

$$\int_M k^2\varphi\bar{\psi} dx = \langle C\varphi, \psi \rangle_H, \quad \varphi, \psi \in H,$$

where $\mathcal{L}(H)$ denotes the space of bounded linear operators from H into itself. Thus the formula (4.6) is equivalent to the operator equation

$$(4.7) \quad w + \mu Bw + (\mu^2 - 1)Cw = 0, \quad w \in H.$$

Further, it is easily verified that B and C are compact operators in $\mathcal{L}(H)$. On the other hand, let $A : \mathbb{C} \rightarrow \mathcal{L}(H)$ be an operator-valued function given by $A(\mu) := \mu B + (\mu^2 - 1)C$. Then it is obvious that $A(\mu)$ is analytic in \mathbb{C} and compact for each $\mu \in \mathbb{C}$. Thus we can apply Corollary 4.1 and the analytic Fredholm theory (see, e.g., [14, Theorem 8.26]) to obtain that $(I + A(\mu))^{-1}$ exists for all $\mu \in \mathbb{C} \setminus S$ where S is a discrete subset of \mathbb{C} with the only accumulating point at infinity. This, together with the equivalence of (BVP) with (4.7), implies the statement of this lemma. \square

Recall from (4.3) and Remark 2.1 that $\sin \theta_n$ are μ -eigenvalues to (BVP) for all $n \in \mathbb{Z}_+$. Since $\theta_n \in (-\pi/2, \pi/2)$ are distinct angles, these μ -eigenvalues must have a finite accumulating point on the real-axis, which contradicts Lemma 4.2. This implies that Case (ii) does not hold.

Finally, the relation $\Gamma^{(1)} = \Gamma^{(2)}$ follows by combining Case (i) and Case (ii). This finishes the proof of Theorem 4.1.

We conclude this section with two remarks.

Remark 4.4. By setting $u = we^{i\alpha x_1}$ with $\alpha \in \mathbb{R}$, the periodic boundary value problem (4.3) can be rewritten as

$$\begin{cases} \Delta u + k^2 u = 0 & \text{in } D, \\ u = 0 & \text{on } \partial D, \\ e^{-i\alpha x_1} u \text{ is } 2\pi\text{-periodic} & \text{with respect to } x_1 \text{ in } D. \end{cases}$$

Multiplying \bar{u} on both sides of the equation and integrating over M , we deduce from the quasi-periodicity of u that

$$0 = \int_M (|\nabla u|^2 - k^2|u|^2) dx.$$

By Poincaré's inequality (see [37, Lemma 3.13]), it follows from the Dirichlet boundary condition of u on $\partial D \cap \bar{M}$ that $0 \geq (C - k^2)\|u\|_{L^2(M)}^2$ for a constant $C > 0$. Hence, $w = e^{-i\alpha x_1} u = 0$ provided $k > 0$ is small enough. Proceeding as in the proof of Lemma 4.2, we can conclude from the analytic Fredholm theory (see, e.g., [14, Theorem 8.26]) that for any $\alpha \in \mathbb{R}$, (4.3) admits only the trivial solution for all $k^2 \in \mathbb{C} \setminus E(\alpha)$ where $E(\alpha)$ is a discrete subset of \mathbb{C} . Therefore, the eigenvalues $\{K_j(\alpha)\}_{j \geq 1}$ are contained in $E(\alpha)$ and thus accumulate only at infinity. Moreover, the associated eigenspace for each eigenvalue $K_j(\alpha)$ is of finite dimension due to the compactness of corresponding operators.

Remark 4.5. In [14, Theorem 5.1], it was proved that a sound-soft scatterer can be uniquely determined by the far-field patterns from a finite number of incident plane waves with a fixed wave number under the assumption that the scatterer is contained in a ball. We note that it is interesting to extend this result to the case of periodic curves. This may require further investigation of properties of the μ -eigenvalues with respect to domains and is thus beyond the scope of this paper. For analogous results with finitely many wave numbers and a fixed incident angle, we refer the reader to [23, Theorem 3.2].

5. Uniqueness with phaseless data. In contrast to the inverse problem with phase information, this section is devoted to uniqueness for recovering the periodic curve from phaseless near-field data (i.e., Theorem 1.1 with phaseless data). We rephrase Theorem 1.1 with phaseless data as follows.

THEOREM 5.1. *Let $\Gamma^{(1)}$ and $\Gamma^{(2)}$ be two periodic curves with Dirichlet boundary conditions. Assume both satisfy Condition (i) or both satisfy Condition (ii). Suppose that the periods of $\Gamma^{(1)}$ and $\Gamma^{(2)}$ are unknown. If the corresponding phaseless total fields satisfy*

$$(5.1) \quad |u_1(x; \theta_n)| = |u_2(x; \theta_n)|, \quad x \in \mathcal{D}, n \in \mathbb{Z}_+,$$

where $\{\theta_n\}_{n=1}^\infty$ are distinct incident angles in $(-\pi/2, \pi/2)$, then $\Gamma^{(1)} = \Gamma^{(2)}$. Here, $\mathcal{D} \subset \Omega$ is a bounded domain.

To prove Theorem 5.1, we will apply Rayleigh expansion (1.4) to show that the phaseless near-field data corresponding to one incident plane wave uniquely determine the total field with phase information except for a finite set of incident angles.

THEOREM 5.2 (phase retrieval). *Let $\Gamma^{(1)}$ and $\Gamma^{(2)}$ be two periodic curves satisfying the conditions in Theorem 5.1. Assume the periods of $\Gamma^{(1)}$ and $\Gamma^{(2)}$ are $L_1 > 0$ and $L_2 > 0$, respectively. Let $u_j(x; \theta)$ ($j = 1, 2$) be the total field for the incident plane wave defined by (1.1) corresponding to the periodic curve $\Gamma^{(j)}$, and let $\theta \in (-\pi/2, \pi/2)$ satisfy $k \sin \theta L_j / \pi \notin \mathbb{Z}$ (i.e., $\alpha L_j / \pi \notin \mathbb{Z}$) for $j = 1, 2$. Suppose the corresponding total fields satisfy*

$$(5.2) \quad |u_1(x; \theta)| = |u_2(x; \theta)|, \quad x \in U_h,$$

for some $h > \max\{x_2 : x \in \Gamma^{(1)} \cup \Gamma^{(2)}\}$. Then $u_1(x; \theta) = u_2(x; \theta)$, $x \in U_h$.

To prove Theorem 5.2, we need several auxiliary lemmas. Let α_n and β_n be defined by (1.5) with some $\theta \in (-\pi/2, \pi/2)$, and let ι be the index for the incident plane wave (see (1.6)).

LEMMA 5.1. *If $\alpha L / \pi \notin \mathbb{Z}$, then $\alpha_n \neq -\alpha_\iota$ for all $n \in \mathbb{Z} \cup \{\iota\}$.*

Proof. We assume to the contrary that $\alpha_n = -\alpha_\iota$ for $n \in \mathbb{Z} \cup \{\iota\}$. Obviously, we have $n \neq \iota$, since otherwise there holds $\alpha_\iota = 0$, which contradicts $\alpha L / \pi \notin \mathbb{Z}$. If $n \in \mathbb{Z}$ and $\alpha + n2\pi / L = -\alpha_\iota = -\alpha$, we can get $\alpha L / \pi = -n \in \mathbb{Z}$, which also contradicts the assumption that $\alpha L / \pi \notin \mathbb{Z}$. \square

In the following, we retain the notation introduced in the proof of Theorem 3.1.

LEMMA 5.2. *Suppose $\Gamma^{(1)}$ and $\Gamma^{(2)}$ are two grating curves with the periods $L_1 > 0$ and $L_2 > 0$, respectively. Assume that $\alpha L_j / \pi \notin \mathbb{Z}$ for $j = 1, 2$. Then the following statements hold.*

(i) *For any fixed $\tilde{m} \in \mathbb{Z}$, if*

$$(5.3) \quad (\alpha_{\tilde{m}}^{(1)} - \alpha_\iota, \beta_{\tilde{m}}^{(1)} - \beta_\iota) = (\alpha_{\tilde{m}}^{(2)} - \alpha_n^{(2)}, \beta_{\tilde{m}}^{(2)} - \overline{\beta_n^{(2)}}),$$

for some $m, n \in \mathbb{Z} \cup \{\iota\}$, then $(\alpha_{\tilde{m}}^{(1)}, \beta_{\tilde{m}}^{(1)}) = (\alpha_m^{(2)}, \beta_m^{(2)})$ and $n = \iota$.

(ii) *For any fixed $\tilde{m} \in \mathbb{Z}$, if*

$$(\alpha_{\tilde{m}}^{(1)} - \alpha_\iota, \beta_{\tilde{m}}^{(1)} - \beta_\iota) = (\alpha_m^{(1)} - \alpha_n^{(1)}, \beta_{\tilde{m}}^{(1)} - \overline{\beta_n^{(1)}}),$$

for some $m, n \in \mathbb{Z} \cup \{\iota\}$, then $(\alpha_{\tilde{m}}^{(1)}, \beta_{\tilde{m}}^{(1)}) = (\alpha_m^{(1)}, \beta_m^{(1)})$ and $n = \iota$.

Proof. We only prove statement (i) since statement (ii) is a consequence of statement (i) for the special case when $\Gamma^{(1)} = \Gamma^{(2)}$.

We consider the following two cases.

Case 1. $\beta_{\bar{m}}^{(1)} \in \mathbb{R}$.

Noting that $\beta_{\bar{m}}^{(1)} - \beta_i > 0$, we conclude from (5.3) that $\beta_m^{(2)}, \beta_n^{(2)} \in \mathbb{R}$. Hence, the points (α_i, β_i) , $(\alpha_{\bar{m}}^{(1)}, \beta_{\bar{m}}^{(1)})$, $(\alpha_m^{(2)}, \beta_m^{(2)})$, and $(\alpha_n^{(2)}, \beta_n^{(2)})$ are all located on the circle $x_1^2 + x_2^2 = k^2$ in the x_1x_2 -plane. From this and the relation (5.3), it follows easily that there holds either

$$(5.4) \quad (\alpha_m^{(2)}, \beta_m^{(2)}) = (\alpha_{\bar{m}}^{(1)}, \beta_{\bar{m}}^{(1)}) \text{ and } (\alpha_n^{(2)}, \beta_n^{(2)}) = (\alpha_i, \beta_i)$$

or

$$(5.5) \quad (\alpha_m^{(2)}, \beta_m^{(2)}) = -(\alpha_i, \beta_i) \text{ and } (\alpha_n^{(2)}, \beta_n^{(2)}) = -(\alpha_{\bar{m}}^{(1)}, \beta_{\bar{m}}^{(1)}).$$

By Lemma 5.1 and the assumption $\alpha L_2/\pi \notin \mathbb{Z}$, the relations in (5.5) cannot be true. Hence, the relations in (5.4) imply the desired result of this lemma.

Case 2. $\beta_{\bar{m}}^{(1)} \notin \mathbb{R}$.

Observing that $\text{Re}(\beta_{\bar{m}}^{(1)} - \beta_i) > 0$ and $\text{Im}(\beta_{\bar{m}}^{(1)} - \beta_i) > 0$, we deduce from (5.3) that $\text{Re}(\beta_m^{(2)} - \beta_n^{(2)}) > 0$ and $\text{Im}(\beta_m^{(2)} - \beta_n^{(2)}) > 0$.

If $\beta_m^{(2)} \notin \mathbb{R}$, then $\beta_m^{(2)}/i \in \mathbb{R}$. This, together with $\text{Re}(\beta_m^{(2)} - \overline{\beta_n^{(2)}}) > 0$, implies $\text{Re}(-\beta_n^{(2)}) > 0$. This is possible only if $n = i$, since $\text{Re}(\beta_n^{(2)}) \geq 0$ for all $n \in \mathbb{Z}$. Again using (5.3), we find $(\alpha_{\bar{m}}^{(1)}, \beta_{\bar{m}}^{(1)}) = (\alpha_m^{(2)}, \beta_m^{(2)})$, which yields the desired result of this lemma.

Now suppose that $\beta_m^{(2)} \in \mathbb{R}$. We shall derive a contradiction as follows. Taking the real and imaginary parts of (5.3) gives $\beta_m^{(2)} = -\beta_i$ and $\beta_n^{(2)} = \beta_{\bar{m}}^{(1)}$. Noting that $(\alpha_m^{(2)})^2 + (\beta_m^{(2)})^2 = k^2 = (\alpha_i)^2 + (\beta_i)^2$, we deduce from $\beta_m^{(2)} = -\beta_i$ that $|\alpha_m^{(2)}| = |\alpha_i|$. Then by $\alpha L_2/\pi \notin \mathbb{Z}$ and Lemma 5.1 we obtain $\alpha_m^{(2)} = \alpha_i$. Inserting this equality into (5.3) gives

$$(5.6) \quad \alpha_{\bar{m}}^{(1)} - \alpha_i = \alpha_m^{(2)} - \alpha_n^{(2)} = \alpha_i - \alpha_n^{(2)}.$$

Similarly, noting that $(\alpha_n^{(2)})^2 + (\beta_n^{(2)})^2 = k^2 = (\alpha_{\bar{m}}^{(1)})^2 + (\beta_{\bar{m}}^{(1)})^2$, we deduce from $\beta_n^{(2)} = \beta_{\bar{m}}^{(1)}$ that $|\alpha_n^{(2)}| = |\alpha_{\bar{m}}^{(1)}|$. If $\alpha_{\bar{m}}^{(1)} = \alpha_n^{(2)}$, then it follows from (5.6) that $\alpha_{\bar{m}}^{(1)} = \alpha_i = \alpha_n^{(2)}$ and thus $\beta_n^{(2)} \in \{\pm\beta_i\} \subset \mathbb{R}$. This contradicts $\beta_n^{(2)} = \beta_{\bar{m}}^{(1)} \notin \mathbb{R}$. If $\alpha_{\bar{m}}^{(1)} = -\alpha_n^{(2)}$, then from (5.6) we deduce $\alpha_i = 0$, which contradicts the assumption $\alpha L_2/\pi \notin \mathbb{Z}$. The proof for Case 2 is complete. \square

With the aid of Lemma 5.2, now we can prove Theorem 5.2.

Proof of Theorem 5.2. Recalling (1.6) and (3.4), we deduce from (5.2) that

$$(5.7) \quad \begin{aligned} & I_1^{(1)}(x)\overline{I_2^{(1)}(x)} + I_2^{(1)}(x)\overline{I_1^{(1)}(x)} + |I_1^{(1)}(x)|^2 + |I_2^{(1)}(x)|^2 \\ & - I_1^{(2)}(x)\overline{I_2^{(2)}(x)} - I_2^{(2)}(x)\overline{I_1^{(2)}(x)} - |I_1^{(2)}(x)|^2 - |I_2^{(2)}(x)|^2 = 0, \quad x \in U_h, \end{aligned}$$

where

$$I_1^{(j)}(x) = \sum_{m \in \mathcal{T}_1^{(j)}} A_m^{(j)} e^{i\alpha_m^{(j)}x_1 + i\beta_m^{(j)}x_2}, \quad I_2^{(j)}(x) = \sum_{n \in \mathcal{T}_2^{(j)}} A_n^{(j)} e^{i\alpha_n^{(j)}x_1 + i\beta_n^{(j)}x_2},$$

with $\mathcal{T}_1^{(j)} := \{n \in \mathbb{Z} : |\alpha_n^{(j)}| > k\}$ and $\mathcal{T}_2^{(j)} := \{n \in \mathbb{Z} \cup \{i\} : |\alpha_n^{(j)}| \leq k\}$, $j = 1, 2$.

The proof can be divided into two steps as follows.

Step 1. We will prove that for any $\tilde{m} \in \mathcal{T}_2^{(1)} \setminus \{i\}$ there holds

$$(5.8) \quad \begin{cases} A_{\tilde{m}}^{(1)} = A_m^{(2)} & \text{if there exists } m \in \mathbb{Z} \text{ such that } \alpha_m^{(2)} = \alpha_{\tilde{m}}^{(1)}, \\ A_{\tilde{m}}^{(1)} = 0 & \text{if } \alpha_m^{(2)} \neq \alpha_{\tilde{m}}^{(1)} \text{ for all } m \in \mathbb{Z}, \end{cases}$$

and for any $\tilde{m} \in \mathcal{T}_2^{(2)} \setminus \{i\}$ there holds

$$(5.9) \quad \begin{cases} A_{\tilde{m}}^{(2)} = A_m^{(1)} & \text{if there exists } m \in \mathbb{Z} \text{ such that } \alpha_m^{(1)} = \alpha_{\tilde{m}}^{(2)}, \\ A_{\tilde{m}}^{(2)} = 0 & \text{if } \alpha_m^{(1)} \neq \alpha_{\tilde{m}}^{(2)} \text{ for all } m \in \mathbb{Z}. \end{cases}$$

First, we deduce (5.8) for $\tilde{m} \in \mathcal{T}_2^{(1)} \setminus \{i\}$. Multiplying (5.7) by $e^{-i(\beta_{\tilde{m}}^{(1)} - \beta_i)x_2}$ we obtain for $x \in U_h$ that

$$(5.10) \quad \begin{aligned} 0 = & \left\{ I_1^{(1)}(x) \overline{I_2^{(1)}(x)} + I_2^{(1)}(x) \overline{I_1^{(1)}(x)} + |I_1^{(1)}(x)|^2 \right\} e^{-i(\beta_{\tilde{m}}^{(1)} - \beta_i)x_2} \\ & + \sum_{(m,n) \in \mathcal{U}_{\tilde{m}}^{(1)}} A_m^{(1)} \overline{A_n^{(1)}} e^{i(\alpha_m^{(1)} - \alpha_n^{(1)})x_1} - \sum_{(m,n) \in \mathcal{U}_{\tilde{m}}^{(2)}} A_m^{(2)} \overline{A_n^{(2)}} e^{i(\alpha_m^{(2)} - \alpha_n^{(2)})x_1} \\ & + \sum_{(m,n) \in (\mathcal{T}_2^{(1)} \times \mathcal{T}_2^{(1)}) \setminus \mathcal{U}_{\tilde{m}}^{(1)}} A_m^{(1)} \overline{A_n^{(1)}} e^{i(\alpha_m^{(1)} - \alpha_n^{(1)})x_1 + i[(\beta_m^{(1)} - \beta_n^{(1)}) - (\beta_{\tilde{m}}^{(1)} - \beta_i)]x_2} \\ & - \left\{ I_1^{(2)}(x) \overline{I_2^{(2)}(x)} + I_2^{(2)}(x) \overline{I_1^{(2)}(x)} + |I_1^{(2)}(x)|^2 \right\} e^{-i(\beta_{\tilde{m}}^{(1)} - \beta_i)x_2} \\ & - \sum_{(m,n) \in (\mathcal{T}_2^{(2)} \times \mathcal{T}_2^{(2)}) \setminus \mathcal{U}_{\tilde{m}}^{(2)}} A_m^{(2)} \overline{A_n^{(2)}} e^{i(\alpha_m^{(2)} - \alpha_n^{(2)})x_1 + i[(\beta_m^{(2)} - \beta_n^{(2)}) - (\beta_{\tilde{m}}^{(1)} - \beta_i)]x_2}, \end{aligned}$$

where $\mathcal{U}_{\tilde{m}}^{(j)} := \{(m, n) \in \mathcal{T}_2^{(j)} \times \mathcal{T}_2^{(j)} : \beta_m^{(j)} - \beta_n^{(j)} = \beta_{\tilde{m}}^{(1)} - \beta_i\}$, $j = 1, 2$. Since $\mathcal{T}_2^{(j)}$ is a finite set, we know that $\mathcal{U}_{\tilde{m}}^{(j)}$ is at most a finite set, $j = 1, 2$. Using $|e^{-i(\beta_{\tilde{m}}^{(1)} - \beta_i)x_2}| = 1$, it follows from Lemma 2.1 (i) that

$$\left| \left\{ I_1^{(j)}(x) \overline{I_2^{(j)}(x)} + I_2^{(j)}(x) \overline{I_1^{(j)}(x)} + |I_1^{(j)}(x)|^2 \right\} e^{-i(\beta_{\tilde{m}}^{(1)} - \beta_i)x_2} \right| \leq C |I_1^{(j)}(x)|, \quad x \in U_h,$$

where $C > 0$ is a constant. Thus, by arguments similar to the proofs of (2.7) and (2.8), we have $|I_1^{(j)}(x)| \rightarrow 0$ as $x_2 \rightarrow +\infty$ and thus

$$\lim_{H \rightarrow +\infty} \frac{1}{H} \int_H^{2H} \left\{ I_1^{(j)}(x) \overline{I_2^{(j)}(x)} + I_2^{(j)}(x) \overline{I_1^{(j)}(x)} + |I_1^{(j)}(x)|^2 \right\} e^{-i(\beta_{\tilde{m}}^{(1)} - \beta_i)x_2} dx_2 = 0$$

uniformly for all $x_1 \in \mathbb{R}$ and $j = 1, 2$. Moreover, it follows easily from Lemma 2.1 (iv) that

$$\lim_{H \rightarrow +\infty} \frac{1}{H} \int_H^{2H} \sum_{(m,n) \in (\mathcal{T}_2^{(j)} \times \mathcal{T}_2^{(j)}) \setminus \mathcal{U}_{\tilde{m}}^{(j)}} A_m^{(j)} \overline{A_n^{(j)}} e^{i(\alpha_m^{(j)} - \alpha_n^{(j)})x_1 + i[(\beta_m^{(j)} - \beta_n^{(j)}) - (\beta_{\tilde{m}}^{(1)} - \beta_i)]x_2} dx_2 = 0$$

uniformly for all $x_1 \in \mathbb{R}$ and $j = 1, 2$. Combining (5.10) and the above formulas, we arrive at

$$\sum_{(m,n) \in \mathcal{U}_{\tilde{m}}^{(1)}} A_m^{(1)} \overline{A_n^{(1)}} e^{i(\alpha_m^{(1)} - \alpha_n^{(1)})x_1} - \sum_{(m,n) \in \mathcal{U}_{\tilde{m}}^{(2)}} A_m^{(2)} \overline{A_n^{(2)}} e^{i(\alpha_m^{(2)} - \alpha_n^{(2)})x_1} = 0, \quad x_1 \in \mathbb{R}.$$

Similarly, multiplying (5.11) by $e^{-i(\alpha_m^{(1)} - \alpha_i)x_1}$, we can employ Lemma 2.1 (iv) to obtain

$$(5.11) \quad \sum_{(m,n) \in \mathcal{V}_{\tilde{m}}^{(1)}} A_m^{(1)} \overline{A_n^{(1)}} - \sum_{(m,n) \in \mathcal{V}_{\tilde{m}}^{(2)}} A_m^{(2)} \overline{A_n^{(2)}} = 0,$$

where $\mathcal{V}_{\tilde{m}}^{(j)} := \{(m, n) \in \mathcal{U}_{\tilde{m}}^{(j)} : \alpha_m^{(j)} - \alpha_n^{(j)} = \alpha_{\tilde{m}}^{(1)} - \alpha_i\}$, $j = 1, 2$. By Lemma 5.2 we have $\mathcal{V}_{\tilde{m}}^{(1)} = \{(\tilde{m}, i)\}$ and $\mathcal{V}_{\tilde{m}}^{(2)} = \{(m, i) : m \in \mathbb{Z} \text{ s.t. } \alpha_m^{(2)} = \alpha_{\tilde{m}}^{(1)}\}$. Thus, noting that $\mathcal{V}_{\tilde{m}}^{(2)}$ is perhaps an empty set and $A_i^{(1)} = 1 = A_i^{(2)}$, we can apply (5.11) to obtain that (5.8) holds for $\tilde{m} \in \mathcal{T}_2^{(1)} \setminus \{i\}$.

Second, by interchanging the roles of $|u_1(x; \theta)|$ and $|u_2(x; \theta)|$, we can employ an argument similar to that above to obtain (5.9) holds for any $\tilde{m} \in \mathcal{T}_2^{(2)} \setminus \{i\}$.

Step 2. We will prove that (5.8) holds for any $\tilde{m} \in \mathcal{T}_1^{(1)}$ and (5.9) holds for any $\tilde{m} \in \mathcal{T}_1^{(2)}$.

By $A_i^{(1)} = A_i^{(2)} = 1$, it follows from (5.7) and the result in Step 1 that

$$(5.12) \quad \begin{aligned} & I_1^{(1)}(x) \overline{I_2^{(1)}(x)} + I_2^{(1)}(x) \overline{I_1^{(1)}(x)} + |I_1^{(1)}(x)|^2 \\ & - I_1^{(2)}(x) \overline{I_2^{(2)}(x)} - I_2^{(2)}(x) \overline{I_1^{(2)}(x)} - |I_1^{(2)}(x)|^2 = 0, \quad x \in U_h. \end{aligned}$$

Let (p_1, q_1) be an element in $\mathcal{B} := \{(1, m) : m \in \mathcal{T}_1^{(1)}\} \cup \{(2, m) : m \in \mathcal{T}_1^{(2)}\}$ such that $|\beta_{q_1}^{(p_1)}| \leq |\beta_m^{(j)}|$ for all $(j, m) \in \mathcal{B}$. Without loss of generality, we assume $p_1 = 1$. Multiplying (5.12) by $e^{-i(\beta_{q_1}^{(1)} - \beta_i)x_2}$ we obtain for $x \in U_h$ that

$$(5.13) \quad \begin{aligned} & \left[I_1^{(1)}(x) e^{-i\beta_{q_1}^{(1)}x_2} \right] \left[\overline{I_2^{(1)}(x)} + \overline{I_1^{(1)}(x)} \right] e^{i\beta_i x_2} + \left[I_2^{(1)}(x) e^{i\beta_i x_2} \right] \left[\overline{I_1^{(1)}(x)} e^{-i\beta_{q_1}^{(1)}x_2} \right] \\ & - \left[I_1^{(2)}(x) e^{-i\beta_{q_1}^{(1)}x_2} \right] \left[\overline{I_2^{(2)}(x)} + \overline{I_1^{(2)}(x)} \right] e^{i\beta_i x_2} + \left[I_2^{(2)}(x) e^{i\beta_i x_2} \right] \left[\overline{I_1^{(2)}(x)} e^{-i\beta_{q_1}^{(1)}x_2} \right] \\ & = 0. \end{aligned}$$

Note that $\beta_m^{(j)} = -\overline{\beta_{q_1}^{(j)}}$ and $|\beta_{q_1}^{(1)}| < |\beta_m^{(j)} - \beta_n^{(j)}|$ for all $m, n \in \mathcal{T}_1^{(j)}$ with $j = 1, 2$. Thus, similarly to the proof of Theorem 3.1, we can apply Lemma 2.1 to obtain that for all $j = 1, 2$ and $x_1 \in \mathbb{R}$,

$$\begin{aligned} \lim_{x_2 \rightarrow +\infty} I_1^{(j)}(x) e^{-i\beta_{q_1}^{(1)}x_2} &= \sum_{m \in \mathcal{T}_1^{(j)} \text{ s.t. } \beta_m^{(j)} = \beta_{q_1}^{(1)}} A_m^{(j)} e^{i\alpha_m^{(j)}x_1}, \\ \lim_{x_2 \rightarrow +\infty} \overline{I_1^{(j)}(x)} e^{-i\beta_{q_1}^{(1)}x_2} &= \sum_{n \in \mathcal{T}_1^{(j)} \text{ s.t. } \beta_n^{(j)} = \beta_{q_1}^{(1)}} \overline{A_n^{(j)}} e^{-i\alpha_n^{(j)}x_1}, \\ \lim_{x_2 \rightarrow +\infty} \left| I_1^{(j)}(x) \right|^2 e^{-i(\beta_{q_1}^{(1)} - \beta_i)x_2} &= 0 \end{aligned}$$

and

$$\begin{aligned} \lim_{H \rightarrow +\infty} \frac{1}{H} \int_H^{2H} I_2^{(j)}(x) e^{i\beta_i x_2} dx_2 &= \sum_{m \in \mathcal{T}_2^{(j)} \text{ s.t. } \beta_m^{(j)} = -\beta_i} A_m^{(j)} e^{i\alpha_m^{(j)}x_1}, \\ \lim_{H \rightarrow +\infty} \frac{1}{H} \int_H^{2H} \overline{I_2^{(j)}(x)} e^{i\beta_i x_2} dx_2 &= \sum_{n \in \mathcal{T}_2^{(j)} \text{ s.t. } \beta_n^{(j)} = \beta_i} \overline{A_n^{(j)}} e^{-i\alpha_n^{(j)}x_1}. \end{aligned}$$

These, together with (5.13), imply for $x_1 \in \mathbb{R}$ that

$$(5.14) \quad \sum_{(m,n) \in \mathcal{U}_{(1,q_1)}^{(1)}} A_m^{(1)} \overline{A_n^{(1)}} e^{i(\alpha_m^{(1)} - \alpha_n^{(1)})x_1} = \sum_{(m,n) \in \mathcal{U}_{(1,q_1)}^{(2)}} A_m^{(2)} \overline{A_n^{(2)}} e^{i(\alpha_m^{(2)} - \alpha_n^{(2)})x_1},$$

where $\mathcal{U}_{q_1}^{(j)} := \{(m, n) \in \mathcal{T}_1^{(j)} \times \mathcal{T}_2^{(j)} : \beta_m^{(j)} = \beta_{q_1}^{(1)}, \beta_n^{(j)} = \beta_i\} \cup \{(m, n) \in \mathcal{T}_2^{(j)} \times \mathcal{T}_1^{(j)} : \beta_m^{(j)} = -\beta_i, \beta_n^{(j)} = \beta_{q_1}^{(1)}\}$ for $j = 1, 2$. It is clear that $\mathcal{U}_{q_1}^{(j)} = \{(m, n) \in (\mathbb{Z} \cup \{i\})^2 : \beta_m^{(j)} - \beta_n^{(j)} = \beta_{q_1}^{(1)} - \beta_i\}$ for $j = 1, 2$. Note that $\mathcal{U}_{q_1}^{(1)}$ and $\mathcal{U}_{q_1}^{(2)}$ are at most finite sets. Then multiplying (5.14) by $e^{-i(\alpha_{q_1}^{(1)} - \alpha_i)x_1}$, we can apply Lemma 2.1 (iv) to obtain

$$(5.15) \quad \sum_{(m,n) \in \mathcal{V}_{q_1}^{(1)}} A_m^{(1)} \overline{A_n^{(1)}} = \sum_{(m,n) \in \mathcal{V}_{q_1}^{(2)}} A_m^{(2)} \overline{A_n^{(2)}},$$

where $\mathcal{V}_{q_1}^{(j)} := \{(m, n) \in \mathcal{U}_{q_1}^{(j)} : \alpha_m^{(j)} - \alpha_n^{(j)} = \alpha_{q_1}^{(1)} - \alpha_i\}$ for $j = 1, 2$. By Lemma 5.2, we have $\mathcal{V}_{q_1}^{(1)} = \{(q_1, i)\}$ and $\mathcal{V}_{q_1}^{(2)} = \{(m, i) : m \in \mathbb{Z} \text{ s.t. } \alpha_m^{(2)} = \alpha_{q_1}^{(1)}\}$. Now we can apply (5.15) and $A_i^{(1)} = 1 = A_i^{(2)}$ to obtain that (5.8) holds for $\tilde{m} = q_1$.

To proceed further, we distinguish between the following two cases.

Case 2.1. there exists $q_2 \in \mathbb{Z}$ such that $\alpha_{q_2}^{(2)} = \alpha_{q_1}^{(1)}$. It is clear that $A_{q_1}^{(1)} = A_{q_2}^{(2)}$ and $q_2 \in \mathcal{T}_1^{(2)}$, and thus we have that (5.9) holds for $\tilde{m} = q_2$. These, together with $A_i^{(1)} = A_i^{(2)} = 1$ and the result in Step 1, imply that $\widehat{I}_2^{(1)}(x) = \widehat{I}_2^{(2)}(x)$ in $x \in U_h$, where

$$\widehat{I}_2^{(j)}(x) = \sum_{n \in \mathcal{T}_2^{(j)} \cup \{q_j\}} A_n^{(j)} e^{i\alpha_n^{(j)}x_1 + i\beta_n^{(j)}x_2}, \quad j = 1, 2.$$

Thus, it follows from (5.2) that

$$\begin{aligned} & \widehat{I}_1^{(1)}(x) \overline{\widehat{I}_2^{(1)}(x)} + \widehat{I}_2^{(1)}(x) \overline{\widehat{I}_1^{(1)}(x)} + |\widehat{I}_1^{(1)}(x)|^2 \\ & - \widehat{I}_1^{(2)}(x) \overline{\widehat{I}_2^{(2)}(x)} - \widehat{I}_2^{(2)}(x) \overline{\widehat{I}_1^{(2)}(x)} - |\widehat{I}_1^{(2)}(x)|^2 = 0, \quad x \in U_h, \end{aligned}$$

where

$$\widehat{I}_1^{(j)}(x) = \sum_{m \in \mathcal{T}_1^{(j)} \setminus \{q_j\}} A_m^{(j)} e^{i\alpha_m^{(j)}x_1 + i\beta_m^{(j)}x_2}, \quad j = 1, 2.$$

Let (p_3, q_3) be an element in $\mathcal{C} := \mathcal{B} \setminus \{(1, q_1), (2, q_2)\}$ s.t. $|\beta_{q_3}^{(p_3)}| \leq |\beta_m^{(j)}|$ for all $(j, m) \in \mathcal{C}$. Then using similar arguments, we can obtain that (5.8) holds for $\tilde{m} = q_3$ if $p_3 = 1$ and (5.9) holds for $\tilde{m} = q_3$ if $p_3 = 2$.

Case 2.2. $\alpha_m^{(2)} \neq \alpha_{q_1}^{(1)}$ for all $m \in \mathbb{Z}$. In this case, $A_{q_1}^{(1)} = 0$. Thus, similarly to Case 2.1, it follows from (5.2) and the result in Step 1 that

$$\begin{aligned} & \widehat{I}_1^{(1)}(x) \overline{\widehat{I}_2^{(1)}(x)} + \widehat{I}_2^{(1)}(x) \overline{\widehat{I}_1^{(1)}(x)} + |\widehat{I}_1^{(1)}(x)|^2 \\ & - \widehat{I}_1^{(2)}(x) \overline{\widehat{I}_2^{(2)}(x)} - \widehat{I}_2^{(2)}(x) \overline{\widehat{I}_1^{(2)}(x)} - |\widehat{I}_1^{(2)}(x)|^2 = 0, \quad x \in U_h, \end{aligned}$$

where $\widehat{I}_1^{(1)}(x)$ is given as in Case 2.1. Let (p_4, q_4) be an element in $\mathcal{E} := \mathcal{B} \setminus \{(1, q_1)\}$ s.t. $|\beta_{q_4}^{(p_4)}| \leq |\beta_m^{(j)}|$ for all $(j, m) \in \mathcal{E}$. Then using similar arguments again, we can obtain that (5.8) holds for $\tilde{m} = q_4$ if $p_4 = 1$ and (5.9) holds for $\tilde{m} = q_4$ if $p_4 = 2$.

For both cases, we can repeat similar arguments to obtain that (5.8) holds for any $\tilde{m} \in \mathcal{T}_1^{(1)}$ and (5.9) holds for any $\tilde{m} \in \mathcal{T}_1^{(2)}$.

Finally, noting that $A_i^{(1)} = A_i^{(2)} = 1$ and combining the results in Step 1 and Step 2, we have $u_1(x; \theta) = u_2(x; \theta)$ for $x \in U_h$. \square

Remark 5.1. The proof for Theorem 5.2 depends only on the Rayleigh expansion (1.4) of the scattered field. Therefore, the phase retrieval result in Theorem 5.2 remains valid under other boundary conditions.

Now we are ready to prove Theorem 5.1.

Proof of Theorem 5.1. For $j = 1, 2$, denote the period of the unknown grating curve $\Gamma^{(j)}$ by $L_j > 0$, and define the set $\mathcal{A} = \{\theta_n : n \in \mathbb{Z}_+ \text{ s.t. } k \sin \theta_n L_j / \pi \notin \mathbb{Z} \text{ for } j = 1, 2\}$, where $\{\theta_n\}_{n=1}^\infty$ are the incident angles from the assumption of Theorem 1.1. By the analyticity of $x \mapsto |u_j(x; \theta)|^2$ in Ω and Theorem 5.2, we have $u_1(x; \theta_n) = u_2(x; \theta_n)$, $x \in U_h$ for any $\theta_n \in \mathcal{A}$. Obviously, $\{\theta \in (-\pi/2, \pi/2) : k \sin \theta L_j / \pi \in \mathbb{Z} \text{ for } j = 1, 2\}$ is a finite set, and thus \mathcal{A} is still an infinite set. Therefore, it follows from Theorem 4.1 that $\Gamma^{(1)} = \Gamma^{(2)}$. \square

Remark 5.2. Assume that the conditions presented in Theorem 5.1 hold true. Assume further that the grating periods L_1 and L_2 are known in advance and $L_1 = L_2$; then the conclusion of Theorem 5.1 can be proved in a very simple way. In fact, let D be the bounded domain defined in subsection 4.1 if $\Gamma^{(1)} \cap \Gamma^{(2)} \neq \emptyset$ or the unbounded periodic strip defined in subsection 4.2 if $\Gamma^{(1)} \cap \Gamma^{(2)} = \emptyset$. Then, due to the analyticity of the total fields and the Dirichlet boundary conditions on $\Gamma^{(1)}$ and $\Gamma^{(2)}$, we can easily deduce from (5.1) that either $\{u_1(x; \theta_n)\}_{n \in \mathbb{Z}_+}$ or $\{u_2(x; \theta_n)\}_{n \in \mathbb{Z}_+}$ satisfy the Helmholtz equation in D with wave number k and vanish on ∂D . This, together with the same arguments as in section 4, gives that $\Gamma^{(1)} = \Gamma^{(2)}$.

6. Conclusion. In this paper, we have established uniqueness results for inverse diffraction grating problems for identifying the period, location, and shape of a periodic curve with Dirichlet boundary condition. Under the a priori smoothness assumption, we proved that the unknown grating curve can be uniquely determined by the near-field data corresponding to infinitely many incident plane waves with different angles at a fixed wave number. For the case when the phase information is not available and the measurement data are taken in a bounded domain above the grating curve, we proved that the phase information can be uniquely determined by phaseless data provided the incident angle θ and the grating period L satisfy the relation $k \sin \theta L / \pi \notin \mathbb{Z}$. Our phase retrieval result (see Theorem 5.2) carries over to other boundary or transmission conditions. However, the proof of Theorem 4.1 for the case $\Gamma^{(1)} \cap \Gamma^{(2)} \neq \emptyset$ does not apply to the Neumann boundary condition, due to the same difficulty for inverse scattering problems by bounded obstacles (see [14, p. 143] for details). In addition, the case that $\Gamma^{(1)} \cap \Gamma^{(2)} = \emptyset$ brings extra difficulties for treating the discreteness of the so-called μ -eigenvalues in a closed waveguide. The uniqueness result with distinct incident angles for recovering penetrable gratings also remains open. Thus it requires new mathematical theory to establish analogues of Theorem 4.1 under other boundary conditions.

Acknowledgment. The authors would like to thank Professor Chunxiong Zheng from Tsinghua University for helpful and stimulating discussions on dispersion relations in a periodic waveguide.

REFERENCES

- [1] T. ABBOUD AND J. C. NEDELEC, *Electromagnetic waves in an inhomogeneous medium*, J. Math. Anal. Appl., 164 (1992), pp. 40–58.

- [2] A. D. AGALTSOV, T. HOHAGE, AND R. G. NOVIKOV, *An iterative approach to monochromatic phaseless inverse scattering*, Inverse Problems, 35 (2019), 024001.
- [3] H. AMMARI, Y. T. CHOW, AND J. ZOU, *Phased and phaseless domain reconstructions in the inverse scattering problem via scattering coefficients*, SIAM J. Appl. Math., 76 (2016), pp. 1000–1030, <https://doi.org/10.1137/15M1043959>.
- [4] T. ARENS, *Scattering by Bi-periodic Layered Media: The Integral Equation Approach*, Habilitationsschrift, KIT, Karlsruhe, 2010.
- [5] T. ARENS AND N. GRINBERG, *A complete factorization method for scattering by periodic surfaces*, Computing, 75 (2005), pp. 111–132.
- [6] G. BAO, *A uniqueness theorem for an inverse problem in periodic diffractive optics*, Inverse Problems, 10 (1994), pp. 335–340.
- [7] G. BAO, P. LI, AND J. LV, *Numerical solution of an inverse diffraction grating problem from phaseless data*, J. Opt. Soc. Amer. A, 30 (2013), pp. 293–299.
- [8] G. BAO, H. ZHANG, AND J. ZOU, *Unique determination of periodic polyhedral structures by scattered electromagnetic fields*, Trans. Amer. Math. Soc., 363 (2011), pp. 4527–4551.
- [9] G. BAO, H. ZHANG, AND J. ZOU, *Unique determination of periodic polyhedral structures by scattered electromagnetic fields II: The resonance case*, Trans. Amer. Math. Soc., 366 (2014), pp. 1333–1361.
- [10] A. S. BONNET-BENDHIA AND F. STARLING, *Guided waves by electromagnetic gratings and non-uniqueness examples for the diffraction problem*, Math. Methods Appl. Sci., 17 (1994), pp. 2305–2338.
- [11] S. N. CHANDLER-WILDE AND J. ELSCHNER, *Variational approach in weighted Sobolev spaces to scattering by unbounded rough surfaces*, SIAM J. Math. Anal., 42 (2010), pp. 2554–2580, <https://doi.org/10.1137/090776111>.
- [12] S. N. CHANDLER-WILDE AND P. MONK, *Existence, uniqueness, and variational methods for scattering by unbounded rough surfaces*, SIAM J. Math. Anal., 37 (2005), pp. 598–618, <https://doi.org/10.1137/040615523>.
- [13] S. N. CHANDLER-WILDE AND B. ZHANG, *A uniqueness result for scattering by infinite rough surfaces*, SIAM J. Appl. Math., 58 (1998), pp. 1774–1790, <https://doi.org/10.1137/S0036139996309722>.
- [14] D. COLTON AND R. KRESS, *Inverse Acoustic and Electromagnetic Scattering Theory*, 4th ed., Springer, New York, 2019.
- [15] D. COLTON AND B. D. SLEEMAN, *Uniqueness theorems for the inverse problem of acoustic scattering*, IMA J. Appl. Math., 31 (1983), pp. 253–259.
- [16] J. ELSCHNER AND G. HU, *Global uniqueness in determining polygonal periodic structures with a minimal number of incident plane waves*, Inverse Problems, 26 (2010), 115002.
- [17] J. ELSCHNER AND G. HU, *Inverse scattering of elastic waves by periodic structures: Uniqueness under the third or fourth kind boundary conditions*, Methods Appl. Anal., 18 (2011), pp. 215–243.
- [18] J. ELSCHNER AND G. HU, *Uniqueness in inverse transmission scattering problems for multilayered obstacles*, Inverse Probl. Imaging, 5 (2011), pp. 793–813.
- [19] J. ELSCHNER, G. SCHMIDT, AND M. YAMAMOTO, *Global uniqueness in determining rectangular periodic structures by scattering data with a single wave number*, J. Inverse Ill-Posed Probl., 11 (2003), pp. 235–244.
- [20] J. ELSCHNER AND M. YAMAMOTO, *An inverse problem in periodic diffractive optics: Reconstruction of Lipschitz grating profiles*, Appl. Anal., 81 (2002), pp. 1307–1328.
- [21] J. ELSCHNER AND M. YAMAMOTO, *Uniqueness results for an inverse periodic transmission problem*, Inverse Probl., 20 (2004), pp. 1841–1852.
- [22] S. FLISS AND P. JOLY, *Solutions of the time-harmonic wave equation in periodic waveguides: Asymptotic behavior and radiation condition*, Arch. Ration. Mech. Anal., 219 (2016), pp. 349–386.
- [23] F. HETTLICH AND A. KIRSCH, *Schiffer’s theorem in inverse scattering theory for periodic structures*, Inverse Probl., 13 (1997), pp. 351–361.
- [24] G. HU AND B. ZHANG, *The linear sampling method for the inverse electromagnetic scattering by a partially coated bi-periodic structure*, Math. Methods Appl. Sci., 34 (2011), pp. 509–519.
- [25] V. ISAKOV, *On uniqueness in the inverse transmission scattering problem*, Comm. Partial Differential Equations, 15 (1990), pp. 1565–1587.
- [26] X. JI, X. LIU, AND B. ZHANG, *Inverse acoustic scattering with phaseless far field data: Uniqueness, phase retrieval, and direct sampling methods*, SIAM J. Imaging Sci., 12 (2019), pp. 1163–1189, <https://doi.org/10.1137/18M1236022>.
- [27] T. KATO, *Perturbation Theory for Linear Operators*, Springer, New York, 1966.

- [28] A. KIRSCH, *Diffraction by periodic structures*, in *Inverse Problems in Mathematical Physics* (Saariselkä, 1992), Lecture Notes in Phys. 422, Springer, Berlin, 1993, pp. 87–102.
- [29] A. KIRSCH, *Uniqueness theorems in inverse scattering theory for periodic structures*, *Inverse Problems*, 10 (1994), pp. 145–152.
- [30] A. KIRSCH AND R. KRESS, *Uniqueness in inverse obstacle scattering*, *Inverse Problems*, 9 (1993), pp. 285–299.
- [31] M. V. KLIBANOV, *Phaseless inverse scattering problems in three dimensions*, *SIAM J. Appl. Math.*, 74 (2014), pp. 392–410, <https://doi.org/10.1137/130926250>.
- [32] M. V. KLIBANOV, *A phaseless inverse scattering problem for the 3-D Helmholtz equation*, *Inverse Probl. Imaging*, 11 (2017), pp. 263–276.
- [33] M. V. KLIBANOV AND V. G. ROMANOV, *Uniqueness of a 3-D coefficient inverse scattering problem without the phase information*, *Inverse Probl.*, 33 (2017), 095007.
- [34] M. V. KLIBANOV, N. A. KOSHEV, D.-L. NGUYEN, L. H. NGUYEN, A. BRETTIN, AND V. N. ASTRATOV, *A numerical method to solve a phaseless coefficient inverse problem from a single measurement of experimental data*, *SIAM J. Imaging Sci.*, 11 (2018), pp. 2339–2367, <https://doi.org/10.1137/18M1179560>.
- [35] R. KRESS, *Linear Integral Equations*, 3rd ed., Appl. Math. Sci. 82, Springer, New York, 2014.
- [36] P. KUCHMENT, *Floquet Theory for Partial Differential Equations*, Birkhäuser, Basel, 1993.
- [37] P. MONK, *Finite Element Methods for Maxwell's Equations*, Oxford University Press, New York, 2003.
- [38] R. G. NOVIKOV, *Formulas for phase recovering from phaseless scattering data at fixed frequency*, *Bull. Sci. Math.*, 139 (2015), pp. 923–936.
- [39] M. REED AND B. SIMON, *Methods of Modern Mathematical Physics. IV. Analysis of Operators*, Academic Press, New York, 1978.
- [40] A. V. SOBOLEV AND J. WALTHOE, *Absolute continuity in periodic waveguides*, *Proc. London Math. Soc.*, 85 (2002), pp. 717–741.
- [41] B. STRYCHARZ, *An acoustic scattering problem for periodic, inhomogeneous media*, *Math. Methods Appl. Sci.*, 21 (1998), pp. 969–983.
- [42] B. STRYCHARZ, *Uniqueness in the inverse transmission scattering problem for periodic media*, *Math. Methods Appl. Sci.*, 22 (1999), pp. 753–772.
- [43] A. WILLERS, *The Helmholtz equation in disturbed half-spaces*, *Math. Methods Appl. Sci.*, 9 (1987), pp. 312–323.
- [44] X. XU, B. ZHANG, AND H. ZHANG, *Uniqueness and direct imaging method for inverse scattering by locally rough surfaces with phaseless near-field data*, *SIAM J. Imaging Sci.*, 12 (2019), pp. 119–152, <https://doi.org/10.1137/18M1210204>.
- [45] X. XU, B. ZHANG, AND H. ZHANG, *Uniqueness in inverse scattering problems with phaseless far-field data at a fixed frequency*, *SIAM J. Appl. Math.*, 78 (2018), pp. 1737–1753, <https://doi.org/10.1137/17M1149699>.
- [46] B. ZHANG AND H. ZHANG, *Recovering scattering obstacles by multi-frequency phaseless far-field data*, *J. Comput. Phys.*, 345 (2017), pp. 58–73.
- [47] B. ZHANG AND H. ZHANG, *Fast imaging of scattering obstacles from phaseless far-field measurements at a fixed frequency*, *Inverse Problems*, 34 (2018), 104005.